antibodies -online.com





Goat anti-Human IgA (Heavy Chain) Antibody (FITC) - Preadsorbed



Go to Product page

1 Image

Overview	
Quantity:	1 mg
Target:	IgA
Binding Specificity:	Heavy Chain
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	FITC
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)
Product Details	
Immunogen:	Immunogen: Human IgA alpha heavy chain
Isotype:	IgG
Characteristics:	Concentration Definition: by UV absorbance at 280 nm
Purification:	Preadsorption: Solid phase absorption
Labeling Ratio:	2.1
Target Details	
Target:	IgA
Target: Abstract:	IgA IgA Products

Target Details

Background:

Synonyms: goat anti-Human IgA (alpha chain) fluorescein conjugated Antibody, goat anti-Human IgA alpha Antibody FITC conjugation

Background: Anti-Human IgA Fluorescein Antibody generated in goat detects immunoglobulin A (alpha chain) from human. Immunoglobulin A (IgA) is an antibody that plays a critical role in mucosal immunity. IgA has two subclasses (IgA1 and IgA2) and can exist in a dimeric form called secretory IgA (sIgA). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Application Details

Application Notes:

Application Note: Anti-Human IgA Fluorescein Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

FLISA Dilution: 1:10,000 - 1:50,000

Flow Cytometry Dilution: 1:500 - 1:2,500

IF Microscopy Dilution: 1:1,000 - 1:5,000

Comment:

Excitation/Emission wavelength: 494 nm/514 nm

Restrictions:

For Research Use only


Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 1.0 mL Reconstitution Buffer: Restore with deionized water (or equivalent)
	Reconstitution burier. Restore with delonized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
	Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.

Handling

Handling Advice:	Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Expiry Date:	12 months

Images



Dot Blot

Image 1. Dot Blot of Anti-HUMAN IgA (alpha chain) (GOAT) Antibody Fluorescein Conjugated. Lane 1: 100ng. Lane 2: 33.3ng. Lane 3: 11.1ng. Lane 4: 3.70ng. Lane 5: 1.23ng. Secondary Antibody: Gt-a-Hu IgA at 1µg/mL. Blocking Buffer: BlockOut for 30 min at RT.