# antibodies .- online.com





# Rabbit anti-Goat IgG (F(ab')2 Region) Antibody (HRP)





Go to Product page

## Overview

Quantity:	1 mg
Target:	IgG
Binding Specificity:	F(ab')2 Region
Reactivity:	Goat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	HRP
Application:	ELISA

## **Product Details**

Immunogen:	Goat IgG fragment
Isotype:	IgG
Specificity:	IgG F(ab')2
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

## Target Details

Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody


## **Application Details**

Application Notes:	This product has been assayed against 1.0 µg of Goat lgG in a standard capture ELISA using
	ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at
	room temperature. A working dilution of 1:8,000 to 1:62,000 of the reconstitution concentration
	is suggested for this product.
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Reconstitution:	Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	Gentamicin sulfate
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	4 °C

#### **Images**



#### **ELISA**

**Image 1.** ELISA results of purified Rabbit Anti-Goat IgG F(ab')2 Antibody Peroxidase Conjugated tested against purified Goat IgG F(ab')2. Each well was coated in duplicate with 1.0  $\mu$ g of Goat IgG F(ab')2. The starting dilution of antibody was  $5\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB-1000.