

## Datasheet for ABIN101310

## Goat anti-Guinea Pig IgG (F(ab')2 Region) Antibody (FITC) - Preadsorbed



Go to Product page

_					
	W	0	rv	10	W

Quantity:	2 mg	
Target:	IgG	
Binding Specificity:	F(ab')2 Region	
Reactivity:	Guinea Pig	
Host:	Goat	
Clonality:	Polyclonal	
Conjugate:	FITC	
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)	
Product Details		
Immunogen:	Immunogen: Guinea Pig IgG F(ab')2 fragment	
Isotype:	IgG	
Isotype: Specificity:	IgG IgG F(ab')2	
Specificity:	IgG F(ab')2	
Specificity: Characteristics:	IgG F(ab')2  Concentration Definition: by UV absorbance at 280 nm	
Specificity: Characteristics: Purification:	IgG F(ab')2  Concentration Definition: by UV absorbance at 280 nm  Preadsorption: Solid phase absorption	
Specificity: Characteristics: Purification: Labeling Ratio:	IgG F(ab')2  Concentration Definition: by UV absorbance at 280 nm  Preadsorption: Solid phase absorption	

## Target Details

Target Type:	Antibody		
Background:	Synonyms: goat Anti-Guinea Pig IgG F(ab')2 Antibody Fluorescein Conjugated, goat Anti-Guine		
	Pig IgG Fab2 Fragment Antibody FITC Conjugated		
	Background: Anti-Guinea Pig IgG F(ab')2 Fluorescein Antibody generated in goat is a proteolytic		
	fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin		
	under controlled conditions of temperature, time and pH . F(ab')2 Molecules lack the Fc portion		
	of IgG and therefore receptors that bind Guinea Pig IgG F(c) will not bind Guinea Pig IgG		
	F(ab')2 Molecules. Secondary Antibodies are available in a variety of formats and conjugate		
	types. When choosing a secondary antibody product, consideration must be given to species		
	and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-		
	reactivity, and host-species source and fragment composition.		
Application Details			
Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence		
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for		
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.		
	FLISA Dilution: 1:10,000 - 1:50,000		
	Flow Cytometry Dilution: 1:500 - 1:2,500		
	IF Microscopy Dilution: 1:1,000 - 1:5,000		
Comment:	Excitation/Emission wavelength: 494 nm/514 nm		
Restrictions:	For Research Use only		
Handling			
Format:	Lyophilized		
Reconstitution:	Reconstitution Volume: 1.0 mL		
	Reconstitution Buffer: Restore with deionized water (or equivalent)		
Concentration:	2.0 mg/mL		
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2		
	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free		
	Preservative: 0.01 % (w/v) Sodium Azide		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		

## Handling

	should be handled by trained staff only.	
Handling Advice:	Product is photosensitive and should be protected from light.	
Storage:	RT,4 °C,-20 °C	
Expiry Date:	12 months	