

Datasheet for ABIN101319

Rabbit anti-Guinea Pig IgG (F(ab')₂ Region) Antibody (Biotin) - Preadsorbed



[Go to Product page](#)

Overview

Quantity:	2 mg
Target:	IgG
Binding Specificity:	F(ab') ₂ Region
Reactivity:	Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

Product Details

Immunogen:	Immunogen: Guinea Pig IgG F(ab') ₂ fragment
Isotype:	IgG
Specificity:	IgG F(ab') ₂
Characteristics:	Concentration Definition: by UV absorbance at 280 nm
Purification:	Preadsorption: immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads

Target Details

Target:	IgG
Abstract:	IgG Products

Target Details

Target Type:	Antibody
Background:	<p>Synonyms: Rabbit Anti-Guinea Pig IgG F(ab')₂ Biotin Conjugated Antibody, Rabbit Anti Guinea Pig IgG F(ab')₂ Antibody Biotin Conjugated, Rabbit Anti Guinea Pig IgG Fab₂ Fragment Antibody Biotin conjugation</p> <p>Background: Anti-Guinea Pig IgG F(ab')₂ Biotin Antibody generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH . F(ab')₂ Molecules lack the Fc portion of IgG and therefore receptors that bind Guinea Pig IgG F(c) will not bind Guinea Pig IgG F(ab')₂ Molecules. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.</p>

Application Details

Application Notes:	<p>Immunohistochemistry Dilution: 1:1,000 - 1:5,000</p> <p>Application Note: This product has been assayed against 1.0 µg of Guinea Pig IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:8,000 to 1:34,000 of the reconstitution concentration is suggested for this product.</p> <p>ELISA Dilution: 1:20,000 - 1:100,000</p> <p>Western Blot Dilution: 1:2,000 - 1:10,000</p>
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	<p>Reconstitution Volume: 1.0 mL</p> <p>Reconstitution Buffer: Restore with deionized water (or equivalent)</p>
Concentration:	2.0 mg/mL
Buffer:	<p>Buffer: 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2</p> <p>Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</p> <p>Preservative: 0.01 % (w/v) Thimerosal</p>
Preservative:	Thimerosal (Merthiolate)

Handling

Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Aliquot to Avoid repeated freezing and thawing.
Storage:	RT,4 °C,-20 °C
Expiry Date:	12 months