

Datasheet for ABIN101798

Sheep anti-Mouse IgG (Heavy & Light Chain) Antibody (Alkaline Phosphatase (AP)) - Preadsorbed



Go to Product pag

1 Image

| Overview | |
|---|--|
| Quantity: | 1 mg |
| Target: | IgG |
| Binding Specificity: | Heavy & Light Chain |
| Reactivity: | Mouse |
| Host: | Sheep |
| Clonality: | Polyclonal |
| Conjugate: | Alkaline Phosphatase (AP) |
| Application: | ELISA, Immunohistochemistry (IHC), Western Blotting (WB) |
| Product Details | |
| Immunogen: | Immunogen: Mouse IgG whole molecule |
| | |
| Isotype: | IgG |
| Isotype: Specificity: | IgG IgG (H&L) |
| | |
| Specificity: | IgG (H&L) |
| Specificity: Characteristics: | IgG (H&L) Concentration Definition: by UV absorbance at 280 nm |
| Specificity: Characteristics: Purification: | IgG (H&L) Concentration Definition: by UV absorbance at 280 nm Preadsorption: Solid phase absorption |
| Specificity: Characteristics: Purification: Sterility: | IgG (H&L) Concentration Definition: by UV absorbance at 280 nm Preadsorption: Solid phase absorption |

Target Details

| Target Type: | Antibody |
|---------------------|--|
| Background: | Synonyms: sheep anti-Mouse IgG Alkaline Phosphatase Conjugated Antibody, sheep anti- |
| | Mouse IgG Antibody Alkaline Phosphatase Conjugation |
| | Background: Anti-Mouse IgG Antibody generated in sheep detects reactivity to Mouse IgG. |
| | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G |
| | constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as |
| | well as fungi and facilitates their destruction or neutralization via agglutination (and thereby |
| | immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. |
| | The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor |
| | proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy |
| | and Light chains of the antibody molecule are present. Secondary Antibodies are available in a |
| | variety of formats and conjugate types. When choosing a secondary antibody product, |
| | consideration must be given to species and immunoglobulin specificity, conjugate type, |
| | fragment and chain specificity, level of cross-reactivity, and host-species source and fragment |
| | composition. |
| | |
| Application Datails | |

Application Details

| Application | Notos: |
|-------------|--------|
| ADDIICATION | Notes. |

Immunohistochemistry Dilution: 1:200 - 1:1,000

Application Note: This product has been assayed against 1.0 µg of Mouse IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concentration is suggested for this product.

ELISA Dilution: 1:10,000 - 1:50,000 Western Blot Dilution: 1:500 - 1:2,500

Restrictions:


For Research Use only

Handling

| Format: | Liquid |
|----------------|--|
| Concentration: | 1.0 mg/mL |
| Buffer: | Buffer: 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50 % (v/v) Glycerol, pH 8.0 |
| | Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| | Preservative: 0.01 % (w/v) Sodium Azide |

Handling

| Preservative: | Sodium azide |
|--------------------|--|
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Handling Advice: | Do not freeze! Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. Do not add Sodium azide. Dilute only prior to immediate use Each reagent is stable for the period shown on the bottle label if stored as directed. |
| Storage: | 4 °C |
| Expiry Date: | 12 months |
| Images | |



Dot Blot

Image 1. Dot Blot of Sheep anti-Mouse IgG Antibody Alkaline Phosphatase Conjugated. Antigen: Mouse IgG. Load: Lane 1 - 200 ng Lane 2 - 66.67 ng Lane 3 - 22.22 ng Lane 4 - 7.41 ng Lane 5 - 2.47 ng. Primary antibody: none. Secondary antibody: Sheep anti-Mouse IgG Antibody Alkaline Phosphatase Conjugated at 1:1,000 for 60 min at RT. Block: ABIN925618 for 60 min at RT. Reaction visualized using alkaline phosphatase substrate (NBT-100) for 30 seconds at RT.