antibodies -online.com





Goat anti-Pig IgG (Heavy & Light Chain) Antibody (FITC) - Preadsorbed



Go to Product pag

2 Images

Overview

Quantity:	2 mg
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Pig
Host:	Goat
Clonality:	Polyclonal
Conjugate:	FITC
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)
Product Details	
Immunogen:	Immunogen: Swine IgG whole molecule
Immunogen: Isotype:	Immunogen: Swine IgG whole molecule IgG
Isotype:	IgG
Isotype: Specificity:	IgG IgG (H&L)
Isotype: Specificity: Characteristics:	IgG IgG (H&L) Concentration Definition: by UV absorbance at 280 nm
Isotype: Specificity: Characteristics: Purification:	lgG lgG (H&L) Concentration Definition: by UV absorbance at 280 nm Preadsorption: immunoaffinity chromatography using Swine lgG coupled to agarose beads
Isotype: Specificity: Characteristics: Purification: Labeling Ratio:	lgG lgG (H&L) Concentration Definition: by UV absorbance at 280 nm Preadsorption: immunoaffinity chromatography using Swine lgG coupled to agarose beads
Isotype: Specificity: Characteristics: Purification: Labeling Ratio: Target Details	IgG (H&L) Concentration Definition: by UV absorbance at 280 nm Preadsorption: immunoaffinity chromatography using Swine IgG coupled to agarose beads 3.64


Target Details

Target Type:	Antibody
Background:	Synonyms: goat Anti-Swine IgG fluorescein Conjugated Antibody, goat Anti-Swine IgG Antibody
	FITC Conjugation
	Background: Anti-Swine IgG (H&L) Fluorescein Antibody generated in goat detects reactivity to
	swine IgG. Secreted as part of the adaptive immune response by plasma B cells,
	immunoglobulin G constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to
	viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via
	agglutination (and thereby immobilizing them), activation of the compliment cascade, and
	opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region,
	recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the
	epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present.
	Secondary Antibodies are available in a variety of formats and conjugate types. When choosing
	a secondary antibody product, consideration must be given to species and immunoglobulin
	specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-
	species source and fragment composition. This Anti-Swine IgG Antibody is conjugated to
	Fluorescein.
Application Details	
Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	FLISA Dilution: 1:10,000 - 1:50,000
	Flow Cytometry Dilution: 1:500 - 1:2,500
	IF Microscopy Dilution: 1:1,000 - 1:5,000
Comment:	Excitation/Emission wavelength: 494 nm/514 nm
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Paconetitution:	Reconstitution Volume: 1.0 mL
Reconstitution:	Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	2.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Handling

	Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Expiry Date:	12 months

Images



Fluorescene Western

Image 1.




Image 2.