ANTIBODIES ONLINE

Datasheet for ABIN102025 Sheep anti-Rabbit IgG (Heavy & Light Chain) Antibody (Texas Red (TR)) - Preadsorbed



Image

Overview

1

Quantity:	2 mg
Target:	lgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rabbit
Host:	Sheep
Clonality:	Polyclonal
Conjugate:	Texas Red (TR)
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)

Product Details

Immunogen:	Immunogen: Rabbit IgG whole molecule
lsotype:	IgG
Specificity:	IgG (H&L)
Characteristics:	Concentration Definition: by UV absorbance at 280 nm
Purification:	Preadsorption: Solid phase absorption
Labeling Ratio:	4.7
Taurant Dataila	
Target Details	

Target:	lgG
Abstract:	IgG Products

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN102025 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target Type:	Antibody
Background:	Synonyms: Sheep Anti Rabbit IgG Antibody Texas Red™ Conjugate, Sheep Anti-Rabbit IgG
	Texas Red™ Conjugated Antibody
	Background: Anti-Rabbit IgG Antibody Texas Red™ generated in sheep detects rabbit IgG.
	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G
	constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as
	well as fungi and facilitates their destruction or neutralization via agglutination (and thereby
	immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis.
	The whole IgG molecule possesses both the $F(c)$ region, recognized by high-affinity Fc receptor
	proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and
	light chains of the antibody molecule are present. Secondary Antibodies are available in a
	variety of formats and conjugate types. When choosing a secondary antibody product,
	consideration must be given to species and immunoglobulin specificity, conjugate type,
	fragment and chain specificity, level of cross-reactivity, and host-species source and fragment
	composition. This Anti-Rabbit IgG (H&L) is conjugated to Texas Red™.

Application Details

Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	FLISA Dilution: 1:10,000 - 1:50,000
	Flow Cytometry Dilution: 1:500 - 1:2,500
	IF Microscopy Dilution: 1:1,000 - 1:5,000
Comment:	Texas Red™ is a registered trademark of Molecular Probes Inc.
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 1.0 mL Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	2.2 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN102025 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Handling

	Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western Blot of Anti-Rabbit IgG (H&L) (SHEEP) Antibody (Min X Hu, Gt, Ms Serum Proteins) . Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule . Lane 2: Rabbit IgG F(ab) Fragment . Lane 3: Rabbit IgG F(c) Fragment . Lane 4: Rabbit IgM Whole Molecule . Lane 5: Normal Rabbit Serum . All samples were reduced. Load: 50 ng per Iane. Block: ABIN925618 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG (H&L) (SHEEP) Antibody (Min X Hu, Gt, Ms Serum Proteins) 1:3,000 for 60 min at RT. Secondary antibody: Anti-Sheep IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in ABIN925618 for 30 min at RT. Predicted/Obsevered Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.