

Datasheet for ABIN102027

Sheep anti-Rabbit IgG (Heavy & Light Chain) Antibody - Preadsorbed



Go to Product pag

1 Image

| Overview | |
|----------------------|--|
| Quantity: | 1 mg |
| Target: | IgG |
| Binding Specificity: | Heavy & Light Chain |
| Reactivity: | Rabbit |
| Host: | Sheep |
| Clonality: | Polyclonal |
| Application: | ELISA, Immunohistochemistry (IHC), Western Blotting (WB) |
| Product Details | |
| Immunogen: | Immunogen: Rabbit IgG whole molecule |
| Isotype: | IgG |
| Specificity: | IgG (H&L) |
| Characteristics: | Concentration Definition: by UV absorbance at 280 nm |
| Purification: | Preadsorption: Solid phase absorption |
| Sterility: | Sterile filtered |
| Target Details | |
| Target: | IgG |
| Abstract: | IgG Products |
| Target Type: | Antibody |

Target Details

Background:

Synonyms: Sheep Anti-Rabbit IgG Antibody Pre-Adsorbed, Sheep Anti-Rabbit IgG (H&L)

Antibody

Background: Anti-Rabbit IgG (H&L) generated in sheep detects rabbit Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75 % of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Application Details

Application Notes:

Immunohistochemistry Dilution: 1:1,000 - 1:5,000

Application Note: Anti-Rabbit IgG antibody is suitable for use in ELISA, immunohistochemistry,

and western blot. Specific conditions for reactivity should be optimized by the end user.

ELISA Dilution: 1:20,000 - 1:100,000


Western Blot Dilution: 1:2,000 - 1:10,000

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|---|
| Concentration: | 1.0 mg/mL |
| Buffer: | Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: None Preservative: 0.01 % (w/v) Sodium Azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Expiry Date: | 12 months |



Western Blotting

Image 1. Western Blot of Anti-Rabbit IgG (H&L) (SHEEP) Antibody (Min X Hu, Gt, Ms Serum Proteins). Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule. Lane 2: Rabbit IgG F(ab) Fragment. Lane 3: Rabbit IgG F(c) Fragment. Lane 4: Rabbit IgM Whole Molecule. Lane 5: Normal Rabbit Serum. All samples were reduced. Load: 50 ng per lane. Block: ABIN925618 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG (H&L) (SHEEP) Antibody (Min X Hu, Gt, Ms Serum Proteins) 1:3,000 for 60 min at RT. Secondary antibody: Anti-Sheep IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in ABIN925618 for 30 min at RT. Predicted/Obsevered Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.