

Datasheet for ABIN102085 Goat anti-Rabbit IgG (Fc Region) Antibody (TRITC) -Preadsorbed



2 Images

Overview

Quantity:	2 mg
Target:	lgG
Binding Specificity:	Fc Region
Reactivity:	Rabbit
Host:	Goat
Clonality:	Polyclonal
Conjugate:	TRITC
Application:	Flow Cytometry (FACS), FLISA, Fluorescence Microscopy (FM)

Product Details

Immunogen:	Immunogen: Rabbit IgG F(c) fragment				
lsotype:	lgG				
Specificity:	lgG F(ab')2				
Characteristics:	Concentration Definition: by UV absorbance at 280 nm				
Purification:	Preadsorption: Solid phase absorption				
Labeling Ratio:	3.0				
Target Details					
Target:	IgG				
Abstract:	IgG Products				

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN102085 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Target Type:	Antibody
Background:	Synonyms: Goat anti-Rabbit IgG F(c) Antibody Rhodamine Conjugation, Goat anti-Rabbit IgG Fc
	fragment Antibody Rhodamine Conjugation, Goat anti-Rabbit IgG F(c) Rhodamine Conjugated
	Antibody
	Background: Anti-Rabbit IgG F(c) Rhodamine antibody generated in goat is a proteolytic
	fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain
	under controlled conditions of temperature, time and pH . Receptors bind the Fc portion of
	rabbit IgG and often this fragment is removed from immunoglobulins to minimize receptor
	binding and lower background reactivity.

Application Details

Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	FLISA Dilution: 1:10,000 - 1:50,000
	Flow Cytometry Dilution: 1:500 - 1:2,500
	IF Microscopy Dilution: 1:1,000 - 1:5,000
Restrictions:	For Research Use only

Handling


Format:	Lyophilized				
Reconstitution:	Reconstitution Volume: 1.0 mL Reconstitution Buffer: Restore with deionized water (or equivalent)				
Concentration:	2.0 mg/mL				
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free Preservative: 0.01 % (w/v) Sodium Azide				
Preservative:	Sodium azide				
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.				
Handling Advice:	Product is photosensitive and should be protected from light.				
Storage:	RT,4 °C,-20 °C				

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN102085 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Expiry Date:

12 months

Images



Dot Blot

Image 1. Dot Blot results of Goat Anti-Rabbit IgG F(c) Antibody Rhodamine Conjugated. Dots are Rabbit IgG at (1) 100ng, (2) 33.3ng, (3) 11.1ng, (4) 3.70ng, (5) 1.23ng. Blocking: ABIN925618 for 30 min at RT. Primary Antibody: none. Secondary Antibody: Goat Anti-Rabbit IgG F(c) FITC at 1µg/mL for 1hr at RT. Imaged with BioRad ChemiDoc, Rhodamine filter.

kDa	1	2	3	4	5
150 -					
100 -					
80 -					
60 -					-
50 -	-				-
40 -					
30 -					
50					
20 -					

Western Blotting

Image 2. Western Blot of Anti-Rabbit IgG F(c) (GOAT) Antibody . Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule . Lane 2: Rabbit IgG F(ab) Fragment . Lane 3: Rabbit IgG F(c) Fragment . Lane 4: Rabbit IgM Whole Molecule . Lane 5: Normal Rabbit Serum . All samples were reduced. Load: 50 ng of IgG, F(ab), IgM and Serum, 100 ng of F(c). Block: ABIN925618 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG F(c) (GOAT) Antibody 1:2,000 for 60 min at RT. Secondary antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in ABIN925618 for 30 min at RT. Predicted/Obsevered Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.