
antibodies -online.com







anti-ATP7A antibody (AA 42-61)





\sim						
	1//	Д	r۱	1	Θ 1	٨

Quantity:	100 μg
Target:	ATP7A
Binding Specificity:	AA 42-61
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Synthetic peptide amino acids 42-61 (cytoplasmic C-terminus) of human Copper-transporting ATPase1
Clone:	S60-4
Isotype:	lgG2b
Specificity:	Detects ~180 kDa in rat brain membrane preparations.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details

Target:	ATP7A		


Target Details

Alternative Name:	ATP7A (ATP7A Products)
Background:	The copper efflux transporters ATP7A and ATP7B sequester intracellular copper into the
	vesicular secretory pathway for export from the cell. ATP7A (also known as Copper-
	transporting ATPase 1) functions as a transmembrane copper-trans locating P-type ATPase
	and plays a vital role in systemic copper absorption in the gut and copper reabsorption in the
	kidney. Polarized epithelial cells such as Madin-Darby canine kidney cells are a physiologically
	relevant model for systemic copper absorption and reabsorption in vivo. Although ATP7A is not
	detectable in most normal tissues, it is expressed in a considerable fraction of many common
	tumor types. Increased expression of ATP7A renders cells resistant to cisplatin and carboplatin
	Mutations in the ATP7A gene result in Menkes disease, which is fatal in early childhood.
	Mutations in the ATP7B gene lead to the autosomal recessive disorder, Wilson disease,
	characterized by neurological symptoms and hepatic damage.
Gene ID:	538
NCBI Accession:	NP_000043
UniProt:	Q04656
Pathways:	Transition Metal Ion Homeostasis, Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	• WB (1:500)
	• ICC/IF (1:100)
	optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN1027714 was sufficient for detection of Copper-transporting ATPase1 in 20 μg
	of rat brain lysate by colorimetric immunoblot analysis using Goat IgG:HRP as the secondary
	antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	-20°C

Images




Immunocytochemistry

Immunocytochemistry/Immunofluorescence 1. **Image** analysis using Mouse Anti-Copper Transporting ATPase 1 Monoclonal Antibody, Clone S60-4 (ABIN1027714). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Copper Transporting **ATPase** Monoclonal Antibody (ABIN1027714) at 1:100 for overnight at 4°C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Copper Transporting ATPase 1 Antibody (D) Composite.



Immunofluorescence (fixed cells)

Image 2. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Copper Transporting ATPase 1 Monoclonal Antibody, Clone S60-4. Tissue: NIH 3T3 (Mouse Fibroblast cell line). Species: Mouse. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Copper Transporting ATPase 1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Endoplasmic Reticulum, Cytoplasm, Golgi Apparatus, Trans-Golgi Network Membrane, Cell Membrane. Magnification:



60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Copper Transporting ATPase 1 Antibody (D) Composite.

Western Blotting

Image 3. Western Blot analysis of Rat Liver showing detection of ~180 kDa Copper Transporting ATPase 1 protein using Mouse Anti-Copper Transporting ATPase 1 Monoclonal Antibody, Clone S60-4 (ABIN1027714). Lane 1: MW Ladder. Lane 2: Rat Liver . Load: 10 ug. Block: 5% Skim Milk powder in TBST. Primary Antibody: Mouse Anti-Copper Transporting ATPase 1 Monoclonal Antibody (ABIN1027714) at 1:500 for 2 hours at RT with shaking. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:4000 for 1 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT. Predicted/Observed Size: ~180 kDa.