

Datasheet for ABIN1043779

## anti-IL17F antibody



[Go to Product page](#)

2 Images

1 Publication

### Overview

Quantity:	100 µg
Target:	IL17F
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

### Product Details

Purpose:	IL-17F Antibody
Immunogen:	Immunogen: Anti-IL-17F (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with mature full length recombinant human IL-17F produced in E.coli followed by hybridoma development. Immunogen Type: Recombinant Protein
Clone:	4H1629-1
Isotype:	IgG1
Cross-Reactivity (Details):	This antibody is specific for human IL-17F protein.
Characteristics:	Synonyms: mouse anti-IL-17F antibody, mouse anti-Interleukin-17F antibody, Cytokine ML-1, Interleukin-24
Purification:	Anti-Human IL-17F (MOUSE) Monoclonal Antibody was purified from concentrated tissue culture supernate by Protein G chromatography followed by extensive dialysis against the buffer stated above.

## Product Details

Sterility: Sterile filtered

## Target Details

Target: IL17F

Alternative Name: IL17F ([IL17F Products](#))

Background: Background: Anti-L-17F recognizes IL-17F (also known as Cytokine ML-1 or Interleukin-24). IL-17F is produced and secreted by CD8+ T cells, NK cells, NKT cells and LT $\alpha$ i cells. The main functions of IL-17F are neutrophil recruitment and immunity to extracellular pathogen. More importantly, IL-17F drives inflammation and auto-immunity. IL-17A and IL-17F are by far the best characterized cytokines of the IL-17 cytokine family. IL-17F dimerizes in a parallel fashion similar to nerve growth factor and other neurotrophins. Its dimerization is critical to fulfill its activity. When secreted by activated T cells, IL-17F can stimulate the production of other cytokines such as IL-6, IL-8 granulocyte colony-stimulating factor and, can stimulate cartilage matrix turnover. Defects in IL17F are the cause of familial candidiasis type 6 (CANDF6). CANDF6 is a rare disorder with altered immune responses and impaired clearance of fungal infections, selective against Candida. Anti-IL-17E cytokine antibody is ideal for investigators involved in Immunology, Signal Transduction research, Cancer and Inflammatory pathologies.

Gene ID: 112744

NCBI Accession: [NP\\_443104](#)

UniProt: [Q96PD4](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Endopeptidase Activity](#)

## Application Details

Application Notes: Immunohistochemistry Dilution: 5  $\mu$ g/mL  
Application Note: Anti-Human IL-17F antibody has been tested for use in IHC and Western Blot. Specific conditions for reactivity should be optimized by the end user.  
Western Blot Dilution: 0.5  $\mu$ g/mL  
ELISA Dilution: 1:10,000-1:50,000  
Other: User Optimized


Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1.0mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: None Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiry Date:	12 months


## Publications

Product cited in:	Conti, Shen, Nayyar, Stocum, Sun, Lindemann, Ho, Hai, Yu, Jung, Filler, Masso-Welch, Edgerton, Gaffen: "Th17 cells and IL-17 receptor signaling are essential for mucosal host defense against oral candidiasis." in: <b>The Journal of experimental medicine</b> , Vol. 206, Issue 2, pp. 299-311, (2009) ( <a href="#">PubMed</a> ).
-------------------	--



Western Blotting

Image 1.



Immunohistochemistry

**Image 2.** Immunohistochemistry of Mouse Anti-IL-17F antibody Tissue: human colon tissue Fixation: formalin-fixed, paraffin-embedded Primary antibody: isotype control (top left) , Mouse Anti-IL-17F antibody (bottom left, right) at 5 ug/ml