

Datasheet for ABIN1043794  
**anti-IL17F antibody (Biotin)**



[Go to Product page](#)

1 Image

1 Publication

## Overview

|              |   |
|--------------|---|
| Quantity:    | 100 µg                                      |
| Target:      | IL17F                                       |
| Reactivity:  | Human                                       |
| Host:        | Mouse                                       |
| Clonality:   | Monoclonal                                  |
| Conjugate:   | This IL17F antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA                |

## Product Details

|                             |  |
|-----------------------------|--|
| Purpose:                    | IL-17F Biotin Conjugated Antibody  |
| Immunogen:                  | Immunogen: Anti-IL-17F (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with mature full length recombinant human IL-17F produced in E.coli followed by hybridoma development.<br>Immunogen Type: Recombinant Protein |
| Clone:                      | 4H450-1  |
| Isotype:                    | IgG1   |
| Cross-Reactivity (Details): | This antibody is specific for human IL-17F protein.  |
| Characteristics:            | Synonyms: mouse anti-IL-17F biotin conjugated antibody, mouse anti-Interleukin-17F biotin conjugated antibody, Cytokine ML-1, Interleukin-24   |
| Purification:               | This product was purified from concentrated tissue culture supernate by Protein G chromatography followed by extensive dialysis against the buffer stated above.   |

## Product Details

Labeling Ratio: 10-20

## Target Details

Target: IL17F

Alternative Name: IL17F ([IL17F Products](#))

Background: Background: Anti-L-17F recognizes IL-17F (also known as Cytokine ML-1 or Interleukin-24). IL-17F is produced and secreted by CD8+ T cells, NK cells, NKT cells and LT $\alpha$ i cells. The main functions of IL-17F are neutrophil recruitment and immunity to extracellular pathogen. More importantly, IL-17F drives inflammation and auto-immunity. IL-17A and IL-17F are by far the best characterized cytokines of the IL-17 cytokine family. IL-17F dimerizes in a parallel fashion similar to nerve growth factor and other neurotrophins. Its dimerization is critical to fulfill its activity. When secreted by activated T cells, IL-17F can stimulate the production of other cytokines such as IL-6, IL-8 granulocyte colony-stimulating factor and, can stimulate cartilage matrix turnover. Defects in IL17F are the cause of familial candidiasis type 6 (CANDF6). CANDF6 is a rare disorder with altered immune responses and impaired clearance of fungal infections, selective against Candida. Anti-IL-17E cytokine antibody is ideal for investigators involved in Immunology, Signal Transduction research, Cancer and Inflammatory pathologies.

Gene ID: 112744

NCBI Accession: [NP\\_443104](#)

UniProt: [Q96PD4](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Endopeptidase Activity](#)

## Application Details

Application Notes: Application Note: This purified antibody has been tested for use in Western Blot. Specific conditions for reactivity should be optimized by the end user.

Western Blot Dilution: 0.2  $\mu$ g/mL

ELISA Dilution: 1:10,000 - 1:50,000

Other: User Optimized

Restrictions: For Research Use only

## Handling


Format: Lyophilized

## Handling

|                    |   |
|--------------------|---|
| Reconstitution:    | Reconstitution Volume: 100 µL<br>Reconstitution Buffer: Restore with deionized water (or equivalent)  |
| Concentration:     | 1.0 mg/mL   |
| Buffer:            | Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2<br>Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free<br>Preservative: 0.01 % (w/v) Sodium Azide   |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.  |
| Storage:           | 4 °C,-20 °C   |
| Storage Comment:   | Store vial at 4° C prior to restoration. Restore with 0.1 mL of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Expiry Date:       | 12 months   |

## Publications

|                   |  |
|-------------------|--|
| Product cited in: | Conti, Shen, Nayyar, Stocum, Sun, Lindemann, Ho, Hai, Yu, Jung, Filler, Masso-Welch, Edgerton, Gaffen: "Th17 cells and IL-17 receptor signaling are essential for mucosal host defense against oral candidiasis." in: <b>The Journal of experimental medicine</b> , Vol. 206, Issue 2, pp. 299-311, (2009) ( <a href="#">PubMed</a> ). |
|-------------------|--|



Western Blotting

**Image 1.** Western Blot of Mouse Anti-IL-17F antibody Lane 1: human full length recombinant IL-17F protein Lane 2: mouse full length recombinant IL-17F protein Lane 3: rat full length recombinant IL-17F protein Load: 20 ng/lane Primary antibody: Anti-IL-17F antibody at 0.2ug/mL for overnight at 4°C