antibodies.com

Datasheet for ABIN1043955 Donkey anti-Guinea Pig IgG (Heavy & Light Chain) Antibody (Biotin) - Preadsorbed



Image

Overview

Quantity:	1 mg
Target:	lgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Guinea Pig
Host:	Donkey
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

Product Details

Immunogen:	Immunogen: Guinea Pig IgG whole molecule
Isotype:	lgG
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Guinea Pig IgG and Guinea Pig Serum.
Cross-Reactivity:	Guinea Pig
Purification:	Preadsorption: immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads
Target Details	
Target:	IgG

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1043955 | 11/30/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Abstract:	IgG Products
Target Type:	Antibody
Background:	Synonyms: Donkey Anti-Guinea Pig IgG Biotin Conjugation, Donkey Anti Guinea Pig IgG Biotin Conjugated
	Background: Anti-Guinea Pig IgG Biotin Antibody generated in donkey detects guinea pig IgG.
	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G
	constitutes 75 % of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as
	well as fungi and facilitates their destruction or neutralization via agglutination (and thereby
	immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis.
	The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor
	proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and
	light chains of the antibody molecule are present. Secondary Antibodies are available in a
	variety of formats and conjugate types. When choosing a secondary antibody product,
	consideration must be given to species and immunoglobulin specificity, conjugate type,
	fragment and chain specificity, level of cross-reactivity, and host-species source and fragment
	composition.

Application Details

Application Notes:	Immunohistochemistry Dilution: 1:1,000 - 1:5,000
	Application Note: This product is designed for immunofluorescence microscopy, fluorescence
	based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for
	multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
	ELISA Dilution: 1:10,000 - 1:50,000
	Western Blot Dilution: 1:2,000 - 1:10,000
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 1.0 mL Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1043955 | 11/30/2023 | Copyright antibodies-online. All rights reserved.

Handling

	Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Centrifuge product if not completely clear after standing at room temperature.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at 4 °C prior to restoration. For extended storage aliquot contents and freeze at -24 °C or below. This product is stable for several weeks at 4 °C as an undiluted liquid.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western Blot of Donkey anti-Guinea Pig IgG Pre-Adsorbed Biotin Conjugated Secondary Antibody. Lane 1: Guinea Pig IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: Biotin conjugated donkey anti-guinea pig IgG 1:1,000 for 60 min at RT. Secondary antibody: Peroxidase Streptavidin secondary antibody at 1:40,000 for 30 min at RT. Block: ABIN925618 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Guinea Pig IgG. Other band(s): None.