

Datasheet for ABIN1044058

Goat anti-Rat IgG (Heavy & Light Chain) Antibody (Texas Red (TR)) - Preadsorbed



Go to Product pag

1 Image

Overview

Quantity:	500 μL
Target:	IgG
Binding Specificity:	Heavy & Light Chain
Reactivity:	Rat
Host:	Goat
Clonality:	Polyclonal
Conjugate:	Texas Red (TR)
Application:	Fluorescence Microscopy (FM)
Product Details	
Immunogen:	Immunogen: Rat IgG whole molecule
Immunogen: Isotype:	Immunogen: Rat IgG whole molecule IgG
Isotype:	IgG
Isotype: Fragment:	lgG F(ab')2 fragment Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat
Isotype: Fragment: Specificity:	IgG F(ab')2 fragment Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG and Rat Serum. This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex

Target Details

Target:	IgG
Abstract:	IgG Products
Target Type:	Antibody
Background:	Synonyms: Goat F(ab')2 Anti-Rat IgG Texas Red™ Conjugated Antibody, Goat Fab2 Anti-Rat IgG
	Antibody Texas Red™ Conjugation
	Background: F(ab')2 Anti-Rat IgG (H&L) Antibody generated in goat detects rat IgG.
	Representing approximately 75 % of serum immunoglobulins, IgG is the most abundant
	antibody isotype found in the circulation. IgG molecules are synthesized and secreted by
	plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types.
	When choosing a secondary antibody product, consideration must be given to species and
	immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-
	reactivity, and host-species source and fragment composition. F(ab')2 Antibody is ideal for
	investigators who routinely perform flow cytometry, immunofluorescence, IHC, and other
	immunoassays. This F(ab')2 Anti-Rat IgG Antibody is conjugated to Texas Red™.

Application Details

Application Notes:	Application Note: This product is designed for immunofluorescence microscopy, fluorescence

based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot

consistency, high titer and specificity.

IF Microscopy Dilution: 1:500-1:2,500

Comment: Texas Red™ is a registered trademark of Molecular Probes Inc.

Restrictions: For Research Use only


Handling

Format:	Lyophilized
Reconstitution:	Reconstitution Volume: 500 µL Reconstitution Buffer: Restore with deionized water (or equivalent)
Concentration:	1.0 mg/mL
Buffer:	Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Handling

	Preservative: 0.01 % (w/v) Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid cycles of freezing and thawing. This vial contains a relatively low volume of reagent (25 μ L). To minimize loss of volume dilute 1:10 by adding 225 μ L of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Product is photosensitive and should be protected from light.
Storage:	RT,4 °C,-20 °C
Storage Comment:	Store vial at -20 °C or below prior to opening. Store the vial at -20 °C or below after dilution.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western Blot of Goat anti-Rat IgG Texas Red Conjugated Antibody. Lane 1: Rat IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Texas Red goat secondary antibody at 1:1,000 for 60 min at RT. Block: ABIN925618 for 30 min at RT. Predicted/Observed size: 28 & 55 kDa, 28 & 55 kDa for Rat IgG. Other band(s): None.