

Datasheet for ABIN1044972

**Sheep Red Blood Cells (100% Washed Pooled Cells)**[Go to Product page](#)**2** Images

## Overview

Quantity:	50 mL
Host:	Sheep
Application:	Haemagglutination (H)

## Product Details

Protein Source:	Red Blood Cells
Characteristics:	Strain: Sheep - Mixed Sex: Mixed
Sterility:	Non-sterile
Components:	Sheep Red Blood Cells (100% Washed Pooled Cells)
Lysate Type:	Normal

## Application Details

Application Notes:	Complement titration, adsorption procedures, HA assays and for the preparation of stroma as particulate reagents.
Comment:	Sheep whole blood is washed to remove the platelet rich plasma, buffy coat layer, and leukocytes (WBC). After processing, the finished product is supplied as 100% red blood cells. Sheep red blood cells are useful for the titration of complement, adsorption procedures, testing for agglutinins/HA assays, and for the preparation of stroma as particulate reagents. Sheep red blood cells are perishable and are collected and processed upon receipt of your order.
Restrictions:	For Research Use only

## Handling

Format: Liquid


Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Storage: 4 °C

Storage Comment: This product MAY be stable for up to one (1) week if properly stored and handled.


Expiry Date: 1 week

## Images



**Image 1.** DNGR signalling promotes phagosomal ROS production. a-b, Confocal images of RAW264.7 cells transfected with empty vector or plasmid encoding C9::C7 or C9(Y7F)::C7 receptors and pulsed with zymosan (a) or dead sRBCs (b) in the presence of Nitroblue tetrazolium (NBT) (Scale bar 10  $\mu$ m). Quantification of ROS+ phagosomes. Data represented as mean ( $\pm$  s.e.m.) (a) or ( $\pm$  s.d.) (b) and are representative of two independent determinations (n = 2). P values determined by one-way ANOVA. c, RAW264.7 stably expressing C9 or C9(Y7F) receptors were pulsed with CellTracker DeepRed (CTDR)-labelled FP-sRBCs for 2 hrs. Percentage of CTDR+ RAW264.7 cells was quantified by flow cytometry. Data represented as mean ( $\pm$  s.d.) and are representative of two independent experiments (n = 2). d, Confocal images of RAW264.7 stably expressing C9 or C9(Y7F) receptors pulsed with dead cells in the presence of NBT for 2 hrs (scale bars 10  $\mu$ m). Image is a representative image of three similar images. e, RT-PCR of NADPH oxidase subunits in HEK293T. Representative of two experiments (n = 2). f, HEK293T cells stably expressing C9::C7 were challenged with zymosan-Oxyburst in the presence or absence of DPI for 1 hr. Oxyburst+ positive phagosomes were quantified across 5 fields of view (n > 100 phagosomes). Data represented as mean ( $\pm$  s.e.m.). P values were calculated by

unpaired parametric test, Mann-Whitney and are representative of two independent experiments (n = 2). g, HEK293T C9::C7 cells were pulsed with zymosan-Ova (left) or transfected with plasmid encoding VENUS-SIINFEKL (right) in the presence or absence of DPI (10  $\mu$ M) for 4 hrs before fixing and adding of OT-I Rag1<sup>-/-</sup> T-cells. IFN- $\gamma$  was assessed by ELISA, plotted as mean ( $\pm$  s.d.) of an experimental triplicate. n.s., not significant, \*P  $\leq$  0.05, \*\*P  $\leq$  0.01. Extended Data Fig 6. PMID: 33349708.



**Image 2.** IgD-only cells have intact germinal center responses but impaired IgG1+ SLPC responses. (A) Splenic (CD19<sup>+</sup>) B cells from WT, IgM<sup>-/-</sup>, and IgD<sup>-/-</sup> mice unimmunized or 5 days after i.p. immunization with 200  $\mu$ L of 10 % SRBCs. (B) Quantification of germinal center (Fashi GL-7hi) cells in (A). (C) Splenocytes from mice in (A). (D) Quantification of CD138<sup>+</sup> IgG1<sup>+</sup> plasma cells in (C). (E) WT (IgM<sup>b+</sup>) and IgM-null (IgD<sup>a+</sup>) germinal center B cells as a percentage of live splenocytes in unimmunized and IgM<sup>+/+</sup> mice 5 days after i.p. immunization with 200  $\mu$ L of 10 % SRBCs. (F) WT (IgG1<sup>b+</sup>) and IgM-null (IgG1<sup>a+</sup>) switched plasma cells (CD138<sup>+</sup> IgG1<sup>+</sup>) as a percentage of live splenocytes in IgM<sup>+/+</sup> mice unimmunized or 5 days after i.p. immunization with 200  $\mu$ L of 10 % SRBCs. (G) Fraction of unswitched NP-specific germinal center cells (CD19<sup>+</sup> Fashi GL-7hi IgM/IgD<sup>+</sup>) from the IgHa locus in the spleens of Balb/c-B6 F1 and IgM<sup>+/+</sup> mice 7-8 days after i.p. immunization with 100  $\mu$ g NP-RSA. (H) Fraction of IgG1+CD138<sup>+</sup> plasma cells from the IgHa locus in Balb/c-B6 F1 and IgM<sup>+/+</sup> mice 7-8 days after i.p. immunization with 100  $\mu$ g NP-RSA. (I) NP-specific IgG1a and IgG1b titers at OD = 0.2 were calculated for the mice in (G-H) by ELISA. The IgG1a to IgG1b titer ratio was calculated for each mouse, and all ratios were normalized such that the average

IgG1a/IgG1b ratio in Balb/c-B6 F1 samples = 1.0. For (A-D), statistics from n = 4 unimmunized mice of each genotype and n = 3 WT, n = 6 IgM<sup>-/-</sup>, and n = 7 IgD<sup>-/-</sup> immunized mice were pooled. For (E-F), n = 5 unimmunized and n = 5 immunized mice are shown. For (G-I), n = 5 Balb/c-B6 F1 mice and n = 3 IgM<sup>+/-</sup> mice are shown. One-way ANOVA with Tukey's multiple comparisons test (B and D), a paired t test (E-F), and Welch's t test (G-I) were used to calculate p values, and mean +SEM is displayed. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001. Figure 7. PMID: 29521626.