

Datasheet for ABIN1048256  
**anti-5HT1D antibody (Cytoplasmic Domain)**

## 2 Images

[Go to Product page](#)

## Overview

|                      |   |
|----------------------|---|
| Quantity:            | 50 µg   |
| Target:              | 5HT1D (HTR1D)   |
| Binding Specificity: | Cytoplasmic Domain  |
| Reactivity:          | Human, Monkey   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Application:         | Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC) |

## Product Details

|                       |   |
|-----------------------|---|
| Brand:                | IHC-plus™   |
| Immunogen:            | <p>Synthetic 16 amino acid peptide from 3rd cytoplasmic domain of human 5HT1D Receptor.</p> <p>Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset (100%), Mouse, Rat, Panda, Bovine, Bat, Horse, Rabbit, Pig, Guinea pig, Turkey, Chicken (94%), Hamster, Elephant, Opossum (88%).</p> <p>Type of Immunogen: Synthetic peptide</p> |
| Specificity:          | Human 5HT1D Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.   |
| Predicted Reactivity: | Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset (100%) Mouse, Rat, Panda, Bovine, Bat, Horse, Rabbit, Pig, Guinea pig, Turkey, Chicken (94%) Hamster, Elephant, Opossum (88%).  |

## Product Details

Purification: Immunoaffinity purified

## Target Details

Target: 5HT1D (HTR1D)

Alternative Name: HTR1D / 5-HT1D Receptor ([HTR1D Products](#))

Background: Name/Gene ID: HTR1D

Subfamily: Serotonin

Family: GPCR

Synonyms: HTR1D, 5-HT1d alpha receptor, 5HT1D Receptor, 5-HT1d-type serotonin receptor, 5-HT1D, 5-HT-1D, 5-HT-1D-alpha, 5-HT1d receptor, HT1d receptor, HTR1DA, HTRL, HT1DA, Serotonin receptor 1D, RDC4, Gpcr14, Serotonin 1d receptor, Serotonin 5-HT-1d receptor

Gene ID: 3352

Pathways: [JAK-STAT Signaling](#)

## Application Details

Application Notes: Approved: IHC, IHC-P (17 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

|               |                       |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

### Handling

---

|         |        |
|---------|--------|
| Format: | Liquid |
|---------|--------|

|                |              |
|----------------|--------------|
| Concentration: | Lot specific |
|----------------|--------------|


|         |                                    |
|---------|------------------------------------|
| Buffer: | PBS, less than 0.1 % sodium azide. |
|---------|------------------------------------|

---

## Handling


|                    |   |
|--------------------|---|
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                      |
| Storage:           | 4 °C,-20 °C   |
| Storage Comment:   | Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles. |
| Expiry Date:       | 12 months   |

## Images



### Immunohistochemistry

**Image 1.** Anti-5HT1D Receptor antibody ABIN1048256 IHC staining of human spinal cord, dorsal root ganglion. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-5HT1D Receptor antibody IHC of human spinal cord, dorsal root ganglion. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.