

Datasheet for ABIN1048263  
**anti-HTR1F antibody (C-Term)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	50 µg
Target:	HTR1F
Binding Specificity:	C-Term
Reactivity:	Human, Rat, Mouse, Monkey, Cow, Rabbit, Dog, Bat, Guinea Pig, Chicken, Hamster
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HTR1F antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC)

## Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 17 amino acid peptide from C-terminus of human 5HT1F Receptor. Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Panda, Dog, Bovine, Bat, Rabbit, Guinea pig, Turkey, Chicken, Platypus (100%), Xenopus, Stickleback, Pufferfish (94%), Opossum, Lizard (88%), Horse (82%).  Type of Immunogen: Synthetic peptide
Specificity:	Human 5HT1F Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except HTR1E (71 %), HTR1D (59 %), HTR1B (59 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan,

## Product Details

---

Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Panda, Dog, Bovine, Bat, Rabbit, Guinea pig, Turkey, Chicken, Platypus (100%) Xenopus, Stickleback, Pufferfish (94%) Opossum, Lizard (88%) Horse (82%).

Purification: Immunoaffinity purified

## Target Details

---

Target: HTR1F

Alternative Name: HTR1F / 5-HT1F Receptor ([HTR1F Products](#))

Background: Name/Gene ID: HTR1F  
Subfamily: Serotonin  
Family: GPCR

Synonyms: HTR1F, 5-HT1F, 5-HT1f receptor, 5HT1F Receptor, 5-HT-1F, 5HT6, HTR1EL, MR77, Serotonin 1f receptor, Serotonin 5-HT-1f receptor, Serotonin receptor 1F

Gene ID: 3355

Pathways: [JAK-STAT Signaling](#)

## Application Details

---

Application Notes: Approved: IHC, IHC-P (2 - 3 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### **Tissue Preparation**

Formalin fixation and embedding in paraffin wax

### **Tissue Sectioning**

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### **Rehydration**

---

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### **Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

## Handling

---


Format: Liquid

---

## Handling


Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images



### Immunohistochemistry

**Image 1.** Anti-HTR1F / 5-HT1F antibody IHC staining of human brain, cortex, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048263 dilution 41335 ug/ml.



### Immunohistochemistry

**Image 2.** Anti-HTR1F / 5-HT1F antibody IHC of human brain, cortex, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 2-3 ug/ml.