

Datasheet for ABIN1048272

anti-Serotonin Receptor 4 antibody (Cytoplasmic Domain)**3** Images[Go to Product page](#)

Overview

Quantity:	50 µg
Target:	Serotonin Receptor 4 (HTR4)
Binding Specificity:	Cytoplasmic Domain
Reactivity:	Human, Cow, Pig, Dog, Horse, Rabbit, Monkey, Bat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	<p>Synthetic 15 amino acid peptide from 3rd cytoplasmic domain of human 5HT4 Receptor.</p> <p>Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Elephant, Panda, Dog, Bat, Bovine, Rabbit, Horse, Pig (100%), Mouse, Rat, Hamster (93%), Guinea pig, Platypus (87%), Lizard (80%).</p> <p>Type of Immunogen: Synthetic peptide</p>
Specificity:	Human 5HT4 Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Elephant, Panda, Dog, Bat, Bovine, Rabbit, Horse, Pig (100%) Mouse, Rat, Hamster (93%) Guinea pig, Platypus (87%) Lizard (80%).

Product Details

Purification: Immunoaffinity purified

Target Details

Target: Serotonin Receptor 4 (HTR4)

Alternative Name: HTR4 / 5-HT4 Receptor ([HTR4 Products](#))

Background: Name/Gene ID: HTR4

Subfamily: Serotonin

Family: GPCR

Synonyms: HTR4, 5-HT4, 5-HT4R, 5-HT-4, 5HT4 receptor, 5-HT4 receptor, 5-hydroxytryptamine 4 receptor, 5-hydroxytryptamine receptor 4, Serotonin 5-HT-4a receptor, Serotonin receptor 4, Cardiac 5-HT4 receptor, Serotonin 4 receptor, Serotonin 5-HT-4 receptor

Gene ID: 3360

Pathways: [JAK-STAT Signaling](#), [cAMP Metabolic Process](#), [S100 Proteins](#)

Application Details

Application Notes: Approved: IHC, IHC-P (15 - 20 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------


Concentration:	Lot specific
----------------	--------------

Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

Handling


Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images




Immunohistochemistry

Image 1. Anti-5HT4 Receptor antibody ABIN1048272 IHC staining of human brain, medulla. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-HTR4 / 5-HT4 Receptor antibody IHC of human Skin, Melanoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Human Kidney: Formalin-Fixed, Paraffin-Embedded (FFPE)