# antibodies -online.com









Go to	D		
	Pron	ויאווו	Dage

_							
0	V	е	r١	/	е	٧	V

Quantity:	50 μg
Target:	BAI3
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Monkey, Chicken, Rabbit, Bat, Dog, Horse, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BAI3 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Brand:	IHC-plus™
Immunogen:	Synthetic 19 amino acid peptide from C-terminus of human BAI3. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Mouse, Rat, Dog, Bat, Bovine, Horse, Rabbit, Opossum, Chicken (100%), Elephant, Panda (95%), Lizard (89%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human BAI3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey,

Panda (95%) Lizard (89%).

Marmoset, Mouse, Rat, Dog, Bat, Bovine, Horse, Rabbit, Opossum, Chicken (100%) Elephant,

# **Product Details** Purification: Immunoaffinity purified **Target Details** Target: BAI3 Alternative Name ADGRB3 / BAI3 (BAI3 Products) Background: Name/Gene ID: ADGRB3 Subfamily: Orphan-B Family: GPCR Synonyms: ADGRB3, KIAA0550, BAI3 Gene ID: 577 **Application Details** Approved: IHC, IHC-P (4 µg/mL) Application Notes: Comment: Target Species of Antibody: Human Assay Procedure: The IHC-pro Immunohistochemistry Protocol **Tissue Preparation** Formalin fixation and embedding in paraffin wax **Tissue Sectioning** Make 4-µm sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-drying oven for 45 minutes at 60°C Deparaffinization Wash slides in 3 changes of xylene - 5 minutes each at room temperature.

# Rehydration

Wash slides in 3 changes of 100% alcohol - 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol - 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

#### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

#### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate - 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

#### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol - 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene - 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only


#### Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

### Handling


	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## **Images**



#### **Immunohistochemistry**

**Image 1.** Anti-BAI3 antibody ABIN1048345 IHC staining of human brain, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



#### **Immunohistochemistry**

**Image 2.** Anti-BAI3 antibody IHC of human Lung, Adenocarcinoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.