

Datasheet for ABIN1048425 anti-CELSR1 antibody (C-Term)





Go to Product page

\sim			
	ve	r\/	٨
\cup	V C	1 V I	٧V

Quantity:	50 μg
Target:	CELSR1
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CELSR1 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™	
Immunogen:	en: Synthetic 19 amino acid peptide from C-terminus of human CELSR1. Percent identity with oth species by BLAST analysis: Human, Gorilla (100%), Gibbon (95%).	
	Type of Immunogen: Synthetic peptide	
Specificity:	Human CELSR1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.	
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla (100%) Gibbon (95%).	
Purification:	Immunoaffinity purified	

Target Details

Target:	CELSR1	
Alternative Name:	CELSR1 (CELSR1 Products)	
Background:	Name/Gene ID: CELSR1	
	Subfamily: Orphan-U Family: GPCR	
	Synonyms: CELSR1, Cadherin family member 9, CDHF9, Flamingo homolog 2, FMI2, HFMI2, Protocadherin flamingo 2	
Gene ID:	9620	
Pathways:	Tube Formation, Asymmetric Protein Localization	

Application Details

Assay Procedure:	The IHC-pro Immunohistochemistry Protocol
Comment:	Target Species of Antibody: Human
Application Notes:	Approved: ELISA, IHC, IHC-P (16 - 32 μg/mL)

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4- μ m sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene - 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol - 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene - 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling


Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months
Imagas	

Images




Immunohistochemistry

Image 1. Anti-CELSR1 antibody IHC staining of human tonsil. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody ABIN1048425 concentration 41372 ug/ml.



Immunohistochemistry

Image 2. Anti-CELSR1 antibody IHC staining of human adrenal. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody ABIN1048425 concentration 41372 ug/ml.



Immunohistochemistry

Image 3. Anti-CELSR1 antibody IHC staining of human testis. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval. Antibody ABIN1048425 concentration 41372 ug/ml.

Please check the product details page for more images. Overall 4 images are available for ABIN1048425.