

Datasheet for ABIN1048440

**anti-Muscarinic Acetylcholine Receptor M2 antibody  
(Cytoplasmic Domain)**[Go to Product page](#)**2** Images

## Overview

|                      |   |
|----------------------|---|
| Quantity:            | 50 µg   |
| Target:              | Muscarinic Acetylcholine Receptor M2 (CHRM2)  |
| Binding Specificity: | Cytoplasmic Domain  |
| Reactivity:          | Human, Mouse, Rat, Cow, Dog, Pig, Monkey, Guinea Pig, Sheep, Rabbit, Horse, Hamster     |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This Muscarinic Acetylcholine Receptor M2 antibody is un-conjugated                     |
| Application:         | Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

## Product Details

|                       |  |
|-----------------------|--|
| Brand:                | IHC-plus™  |
| Immunogen:            | Synthetic 16 amino acid peptide from 3rd cytoplasmic domain of human CHRM2. Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan, Gibbon, Monkey, Marmoset, Mouse, Rat, Sheep, Dog, Bovine, Hamster, Elephant, Panda, Horse, Rabbit, Pig, Guinea pig (100%), Bat, Opossum, Turkey, Chicken (94%), Xenopus (88%), Platypus, Zebrafish, Stickleback (81%).<br><br>Type of Immunogen: Synthetic peptide |
| Specificity:          | Human CHRM2. BLAST analysis of the peptide immunogen showed no homology with other human proteins.   |
| Predicted Reactivity: | Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Orangutan,  |

## Product Details

---

Gibbon, Monkey, Marmoset, Mouse, Rat, Sheep, Dog, Bovine, Hamster, Elephant, Panda, Horse, Rabbit, Pig, Guinea pig (100%) Bat, Opossum, Turkey, Chicken (94%) Xenopus (88%) Platypus, Zebrafish, Stickleback (81%).

Purification: Immunoaffinity purified

## Target Details

---

Target: Muscarinic Acetylcholine Receptor M2 (CHRM2)

Alternative Name: CHRM2 / M2 ([CHRM2 Products](#))

Background: Name/Gene ID: CHRM2  
Subfamily: Acetylcholine  
Family: GPCR

Synonyms: CHRM2, 7TM receptor, HM2, M2 muscarinic receptor, M2-mAChR, Muscarinic receptor M2, Muscarinic M2 receptor

Gene ID: 1129

## Application Details

---

Application Notes: Approved: IHC, IHC-P (6 - 15 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### **Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid


Concentration: Lot specific

---

## Handling


|                    |   |
|--------------------|---|
| Buffer:            | PBS, less than 0.1 % sodium azide.  |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                      |
| Storage:           | 4 °C, -20 °C  |
| Storage Comment:   | Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles. |
| Expiry Date:       | 12 months   |

## Images



### Immunohistochemistry

**Image 1.** Anti-CHRM2 antibody ABIN1048440 IHC staining of human brain, neurons and glia. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-CHRM2 / M2 antibody IHC of human Brain, Glioblastoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.