

Datasheet for ABIN1048464  
**anti-CRHR2 antibody (N-Term)**



[Go to Product page](#)

2 Images

## Overview

|                      |   |
|----------------------|---|
| Quantity:            | 50 µg   |
| Target:              | CRHR2   |
| Binding Specificity: | N-Term  |
| Reactivity:          | Human, Cow, Rabbit, Monkey  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This CRHR2 antibody is un-conjugated  |
| Application:         | Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

## Product Details

|                       |   |
|-----------------------|---|
| Brand:                | IHC-plus™   |
| Immunogen:            | Synthetic 19 amino acid peptide from N-terminal extracellular domain of human CRHR2.<br>Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Elephant, Panda, Bovine, Rabbit (100%), Mouse, Rat, Dog, Horse, Pig (95%), Opossum, Turkey, Lizard, Stickleback, Zebrafish (89%), Chicken (84%).<br><br>Type of Immunogen: Synthetic peptide |
| Specificity:          | Human CRHR2. BLAST analysis of the peptide immunogen showed no homology with other human proteins.  |
| Predicted Reactivity: | Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Elephant, Panda, Bovine, Rabbit (100%) Mouse, Rat, Dog, Horse, Pig (95%) Opossum,  |

## Product Details

---

Turkey, Lizard, Stickleback, Zebrafish (89%) Chicken (84%).

Purification: Immunoaffinity purified

## Target Details

---

Target: CRHR2

Alternative Name: CRHR2 / CRF2 Receptor ([CRHR2 Products](#))

Background: Name/Gene ID: CRHR2  
Subfamily: Corticotropin releasing factor  
Family: GPCR

Synonyms: CRHR2, CRF-R-2, CRF-R2, CRF2R, Crh2 receptor, CRF2 receptor, beta isoform, CRFR2, CRFR-2, CRH-R-2, CRH2R, CRF-RB, CRF2, CRH-R2, HM-CRF

Gene ID: 1395

Pathways: [Negative Regulation of Hormone Secretion](#), [cAMP Metabolic Process](#), [Feeding Behaviour](#)

## Application Details

---

Application Notes: Approved: IHC, IHC-P (8 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### **Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid


Concentration: Lot specific

---

## Handling


|                    |   |
|--------------------|---|
| Buffer:            | PBS, less than 0.1 % sodium azide.  |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                      |
| Storage:           | 4 °C,-20 °C   |
| Storage Comment:   | Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles. |
| Expiry Date:       | 12 months   |

## Images



### Immunohistochemistry

**Image 1.** Anti-CRHR2 antibody ABIN1048464 IHC staining of human brain, pituitary. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-CRHR2 antibody IHC of human brain, pituitary. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.