

Datasheet for ABIN1048487  
**anti-CysLTR1 antibody (Extracellular Domain)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	50 µg
Target:	CysLTR1 (CYSLTR1)
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CysLTR1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 20 amino acid peptide from 3rd extracellular domain of human CYSLTR1 / CYSLT1. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey (100%), Rabbit, Guinea pig (85%), Bovine, Elephant, Horse, Pig (80%).  Type of Immunogen: Synthetic peptide
Specificity:	Human CYSLTR1 / CYSLT1. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except NMU (45 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey (100%) Rabbit, Guinea pig (85%) Bovine, Elephant, Horse, Pig (80%).
Purification:	Immunoaffinity purified

## Target Details

---

Target: CysLTR1 (CYSLTR1)

Alternative Name: CYSLTR1 / CYSLT1 ([CYSLTR1 Products](#))

Background: Name/Gene ID: CYSLTR1  
Subfamily: Chemoattractant  
Family: GPCR

Synonyms: CYSLTR1, Cys-LT1 receptor, CysLT(1), Cys-leukotriene-1 receptor, CYSLT1, CYSLT1R, CYSLTR, HG55, LTD4 receptor, HMTMF81, CysteinyI It1 receptor

Gene ID: 10800

## Application Details

---

Application Notes: Approved: IHC, IHC-P (6 - 14 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### **Tissue Preparation**

Formalin fixation and embedding in paraffin wax

### **Tissue Sectioning**

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### **Rehydration**

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### **Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: Lot specific

Buffer: PBS, less than 0.1 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C


---

## Handling

Storage Comment: Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.


Expiry Date: 12 months

## Images



### Immunohistochemistry

**Image 1.** Anti-CYSLTR1 / CYSLT1 antibody ABIN1048487 IHC staining of human nasal mucosa, allergic rhinitis, eosinophils. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-CYSLTR1 / CYSLT1 antibody IHC of human Lymph Node, Hodgkins Lymphoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.