

Datasheet for ABIN1048611
anti-FZD2 antibody (Extracellular Domain)

2 Images

[Go to Product page](#)

Overview

Quantity:	50 µg
Target:	FZD2
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Rat, Mouse, Horse, Cow, Dog, Hamster, Pig
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	<p>Synthetic 18 amino acid peptide from 3rd extracellular domain of human Frizzled 2. Percent identity with other species by BLAST analysis: Human, Mouse, Rat, Hamster, Elephant, Panda, Dog, Bovine, Horse, Pig (100%), Sheep, Rabbit (94%), Opossum (89%), Turkey, Chicken, Platypus, Lizard, Xenopus (83%).</p> <p>Type of Immunogen: Synthetic peptide</p>
Specificity:	Human Frizzled 2. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except FZD7 (67 %), OR1A2 (56 %), NFATC2 (50 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Mouse, Rat, Hamster, Elephant, Panda, Dog, Bovine, Horse, Pig (100%) Sheep, Rabbit (94%) Opossum (89%) Turkey, Chicken, Platypus, Lizard, Xenopus (83%).

Product Details

Purification: Immunoaffinity purified

Target Details

Target: FZD2

Alternative Name: FSD2 / Frizzled 2 ([FZD2 Products](#))

Background: Name/Gene ID: FZD2
Subfamily: Frizzled
Family: GPCR

Synonyms: FZD2, Dfrizzled2, Fz2, Frizzled family receptor 2, Frizzled-2, HFz2, Frizzled homolog 2, Fz-2, FzE2, Mfz10, Rfz2, Frizzled 2

Gene ID: 2535

Pathways: [WNT Signaling](#)

Application Details

Application Notes: Approved: IHC, IHC-P (5 - 10 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------


Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

Preservative:	Sodium azide
---------------	--------------

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-Frizzled 2 antibody ABIN1048611 IHC staining of human fetal kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Human Kidney: Formalin-Fixed, Paraffin-Embedded (FFPE)