

Datasheet for ABIN1048630
anti-FZD8 antibody (Extracellular Domain)

2 Images

[Go to Product page](#)

Overview

Quantity:	50 µg
Target:	FZD8
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Mouse, Rat, Dog, Cow, Monkey, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 16 amino acid peptide from 3rd extracellular domain of human FZD8 / Frizzled 8. Percent identity with other species by BLAST analysis: Human, Marmoset, Mouse, Rat, Elephant, Panda, Bovine, Dog, Rabbit, Opossum (100%), Lizard (94%), Monkey (81%). Type of Immunogen: Synthetic peptide
Specificity:	Human FZD8 / Frizzled 8. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except INTS6 (50 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Marmoset, Mouse, Rat, Elephant, Panda, Bovine, Dog, Rabbit, Opossum (100%) Lizard (94%) Monkey (81%).
Purification:	Immunoaffinity purified

Target Details

Target:	FZD8
Alternative Name:	FZD8 / Frizzled 8 (FZD8 Products)
Background:	Name/Gene ID: FZD8 Subfamily: Frizzled Family: GPCR Synonyms: FZD8, Frizzled family receptor 8, Frizzled homolog 8, FZ-8, Fz8, Frizzled 8, Frizzled-8, HFZ8
Gene ID:	8325
Pathways:	WNT Signaling

Application Details

Application Notes:	Approved: IHC, IHC-P (8 - 20 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	The IHC-pro Immunohistochemistry Protocol

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes
Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.
Apply a universal protein block – 20 minutes at room temperature.
Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.
Rinse slides in 1X TBST - 1 minute at room temperature.
Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.
Rinse slides 1X TBST – 1 minute at room temperature.
Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.
Rinse slides in 1X TBST - 1 minute at room temperature.
Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.
Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.
Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.
Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.
Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.
Wash slides in 3 changes of xylene – 1 minute each at room temperature.
Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling


Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-FZD8 / Frizzled 8 antibody IHC staining of human brain, cerebellum, purkinje neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048630 dilution 44044 ug/ml.



Immunohistochemistry

Image 2. Anti-FZD8 / Frizzled 8 antibody IHC staining of human fetal kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048630 dilution 44044 ug/ml.