

Datasheet for ABIN1048636

anti-GABBR1 antibody (N-Term)

50 μg





Go to Product page

()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Quantity:

~		
Target:	GABBR1	
Binding Specificity:	N-Term	
Reactivity:	Human, Rat, Mouse, Cow, Monkey, Pig, Hamster, Rabbit	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GABBR1 antibody is un-conjugated	
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Brand:	IHC-plus [™]	
Immunogen:	Synthetic 15 amino acid peptide from N-terminal extracellular domain of human GABBR1. Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Monkey, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Rabbit, Pig, Opossum (100%), Horse (87%), Xenopus, Stickleback, Zebrafish (80%).	
	Type of Immunogen: Synthetic peptide	
Specificity:	Human GABBR1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.	
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Monkey,	
	Mouse, Rat, Hamster, Elephant, Panda, Bovine, Rabbit, Pig, Opossum (100%) Horse (87%)	

Product Details

	Xenopus, Stickleback, Zebrafish (80%).	
Purification:	Immunoaffinity purified	
Target Details		
Target:	GABBR1	
Alternative Name:	GABABR1 / GABBR1 (GABBR1 Products)	
Background:	Name/Gene ID: GABBR1	
	Subfamily: GABA(B)	
	Family: GPCR	
	Synonyms: GABBR1, DJ271M21.1.1, DJ271M21.1.2, GABA(B) receptor, Gaba-b-r1a, GABA-BR1,	
	Gaba-br1a receptor, GABAB, subunit 1c, GABAB1, GABA-B receptor 1, Gaba-b1, Gababr1	
	receptor, GPRC3A, Gabab-receptor subunit 1a, GABABR1, GABBR1-3, GBR1, HGB1a, GABA-B-	
	R1, GB1	
Gene ID:	2550	
Pathways:	Positive Regulation of Peptide Hormone Secretion, cAMP Metabolic Process	
Application Details		
Application Notes:	Approved: IHC, IHC-P (9 - 18 μg/mL)	
Comment:	Target Species of Antibody: Human	
Assay Procedure:	The IHC-pro Immunohistochemistry Protocol	
	Tissue Preparation	
	Formalin fixation and embedding in paraffin wax	
	Tissue Sectioning	
	Make 4-µm sections and place on pre-cleaned and charged microscope slides.	
	Heat in a tissue-drying oven for 45 minutes at 60°C	
	Deparaffinization	
	Wash slides in 3 changes of xylene – 5 minutes each at room temperature.	

Wash slides in 3 changes of 100% alcohol - 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol - 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate - 30 minutes at room temperature.

Wash slides in distilled water - 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only

Handling


Format:

Liquid

Handling


Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-GABBR1 antibody ABIN1048636 IHC staining of human brain, Purkinje neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-GABABR1 / GABBR1 antibody IHC of human Ovary, Carcinoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.