antibodies -online.com





anti-GABBR1 antibody (N-Term)

2 Images



Go to Product page

\sim					
	1//6	r	V I	Θ	Λ

Predicted Reactivity:

Quantity:	50 μg		
Target:	GABBR1		
Binding Specificity:	N-Term		
Reactivity:	Human, Rat, Mouse, Monkey, Cow, Pig, Hamster, Dog, Horse, Bat		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This GABBR1 antibody is un-conjugated		
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))		
Product Details			
Product Details Brand:	IHC-plus™		
	IHC-plus™ Synthetic 15 amino acid peptide from N-terminal extracellular domain of human GABBR1. Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Bat, Horse, Pig (100%), Rabbit, Platypus (93%), Lizard, Xenopus, Stickleback, Zebrafish (87%), Opossum (80%).		
Brand:	Synthetic 15 amino acid peptide from N-terminal extracellular domain of human GABBR1. Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Bat, Horse, Pig (100%),		

Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Gibbon,

Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Bat, Horse, Pig (100%)

Product Details

	Rabbit, Platypus (93%) Lizard, Xenopus, Stickleback, Zebrafish (87%) Opossum (80%).		
Purification:	Immunoaffinity purified		
Target Details			
Target:	GABBR1		
Alternative Name:	GABABR1 / GABBR1 (GABBR1 Products)		
Background:	Name/Gene ID: GABBR1		
	Subfamily: GABA(B)		
	Family: GPCR		
	Synonyms: GABBR1, DJ271M21.1.1, DJ271M21.1.2, GABA(B) receptor, Gaba-b-r1a, GABA-BR1		
	Gaba-br1a receptor, GABAB, subunit 1c, GABAB1, GABA-B receptor 1, Gaba-b1, Gababr1		
	receptor, GPRC3A, Gabab-receptor subunit 1a, GABABR1, GABBR1-3, GBR1, HGB1a, GABA-B-		
	R1, GB1		
Gene ID:	2550		
Pathways:	Positive Regulation of Peptide Hormone Secretion, cAMP Metabolic Process		
Application Details			
Application Notes:	Approved: IHC, IHC-P (7 - 13 μg/mL)		
Comment:	Target Species of Antibody: Human		
Assay Procedure:	The IHC-pro Immunohistochemistry Protocol		
	Tissue Preparation		
	Formalin fixation and embedding in paraffin wax		
	Tissue Sectioning		
	Make 4-µm sections and place on pre-cleaned and charged microscope slides.		
	Heat in a tissue-drying oven for 45 minutes at 60°C		
	Deparaffinization		
	Wash slides in 3 changes of xylene – 5 minutes each at room temperature.		

Wash slides in 3 changes of 100% alcohol - 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water - 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only

Handling


Format:

Liquid

Handling


Concentration:	Lot specific	
Buffer:	PBS, less than 0.1 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C fo up to 1 month. Avoid freeze-thaw cycles.	
Expiry Date:	12 months	

Images



Immunohistochemistry

Image 1. Anti-GABBR1 antibody ABIN1048637 IHC staining of human brain, neurons and glia. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-GABABR1 / GABBR1 antibody IHC of human Brain, Glioblastoma. Immunohistochemistry of formalinfixed, paraffin-embedded tissue after heat-induced antigen retrieval.