.-online.com antibodies

Datasheet for ABIN1048655 anti-GIPR antibody (N-Term)

2 Images



Overview

Quantity:	50 µg
Target:	GIPR
Binding Specificity:	N-Term
Reactivity:	Human, Cow, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GIPR antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 17 amino acid peptide from N-terminal extracellular domain of human GIPR. Percent identity with other species by BLAST analysis: Human, Gorilla, Bovine, Dog, Panda (100%), Horse, Rabbit (94%), Marmoset, Hamster (88%), Rat (82%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human GIPR. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except GHRHR (100 %), TG (41 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Bovine, Dog, Panda (100%) Horse, Rabbit (94%) Marmoset, Hamster (88%) Rat (82%).
Purification:	Immunoaffinity purified

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN1048655 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Target Details

i di got 2 ottino	
Target:	GIPR
Alternative Name:	GIPR / GIP Receptor (GIPR Products)
Background:	Name/Gene ID: GIPR
	Subfamily: Gastric inhibitory polypeptide
	Family: GPCR
	Synonyms: GIPR, GIP receptor, PGQTL2, GIP-R
Gene ID:	2696
Pathways:	Positive Regulation of Peptide Hormone Secretion, cAMP Metabolic Process, Regulation of G-
	Protein Coupled Receptor Protein Signaling
Application Details	
Application Notes:	Approved: IHC, IHC-P (16 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	The IHC-pro Immunohistochemistry Protocol
	Tissue Preparation
	Formalin fixation and embedding in paraffin wax
	Tissue Sectioning
	Make 4-µm sections and place on pre-cleaned and charged microscope slides.
	Heat in a tissue-drying oven for 45 minutes at 60°C
	Deparaffinization
	Wash slides in 3 changes of xylene – 5 minutes each at room temperature.
	Rehydration
	Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.
	Weak alides in Ω abandon of ΩE^{0} alookal. Ω minutes each at room temperature

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN1048655 | 09/12/2023 | Copyright antibodies-online. All rights reserved. Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure. Apply a universal protein block – 20 minutes at room temperature. Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature. Rinse slides in 1X TBST - 1 minute at room temperature. Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature. Rinse slides 1X TBST – 1 minute at room temperature. Apply alkaline phosphatase streptavidin – 30 minutes at room temperature. Rinse slides in 1X TBST - 1 minute at room temperature. Apply alkaline phosphatase streptavidin – 30 minutes at room temperature. May alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble. Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature. Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature. Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature. Wash slides in 3 changes of xylene – 1 minute each at room temperature. Apply coverslip

Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN1048655 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images





Immunohistochemistry

Image 1. Anti-GIPR antibody ABIN1048655 IHC staining of human pancreas. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Immunohistochemistry

Image 2. Anti-GIPR antibody IHC of human pancreas. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN1048655 | 09/12/2023 | Copyright antibodies-online. All rights reserved.