

Datasheet for ABIN1048698

**anti-G Protein-Coupled Receptor 119 antibody (Cytoplasmic Domain)**[Go to Product page](#)**3** Images

## Overview

Quantity:	50 µg
Target:	G Protein-Coupled Receptor 119 (GPR119)
Binding Specificity:	Cytoplasmic Domain
Reactivity:	Human, Bat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This G Protein-Coupled Receptor 119 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 20 amino acid peptide from 3rd cytoplasmic domain of human GPR119. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Bat (100%), Marmoset, Mouse, Hamster, Panda, Dog, Horse (95%), Rat, Elephant, Rabbit (90%), Bovine, Pig (80%).  Type of Immunogen: Synthetic peptide
Specificity:	Human GPR119. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Bat (100%) Marmoset, Mouse, Hamster, Panda, Dog, Horse (95%) Rat, Elephant, Rabbit (90%)

## Product Details

Bovine, Pig (80%).

Purification: Immunoaffinity purified

## Target Details

Target: G Protein-Coupled Receptor 119 (GPR119)

Alternative Name: GPR119 ([GPR119 Products](#))

Background: Name/Gene ID: GPR119  
Subfamily: Lysophospholipid/Lysosphingolipid  
Family: GPCR

Synonyms: GPR119, G-protein coupled receptor 2, GPCR2, HGPCR2, G protein-coupled receptor 119, G-protein coupled receptor 119

Gene ID: 139760

Pathways: [Hormone Transport, Peptide Hormone Metabolism](#)

## Application Details

Application Notes: Approved: IHC, IHC-P (5 µg/mL)

Usage: Proteinase K treatment is required for antigen retrieval.

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### **Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---


Format:	Liquid
---------	--------

---

## Handling


Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images




### Immunohistochemistry

**Image 1.** Anti-GPR119 antibody ABIN1048698 IHC staining of human brain, cortex. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-GPR119 antibody ABIN1048698 IHC staining of human liver. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



#### Immunohistochemistry

**Image 3.** Anti-GPR119 antibody IHC of human brain, cortex. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.