



Datasheet for ABIN1048704
anti-GPR124 antibody (N-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	50 µg
Target:	GPR124
Binding Specificity:	N-Term
Reactivity:	Human, Bat, Dog, Horse, Monkey, Pig, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 17 amino acid peptide from N-terminal extracellular domain of human GPR124. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Marmoset, Dog, Bat, Elephant, Panda, Horse, Rabbit, Pig, Opossum (100%), Mouse, Rat, Hamster, Turkey, Chicken, Xenopus (94%), Monkey, Bovine, Platypus, Lizard (82%). Type of Immunogen: Synthetic peptide
Specificity:	Human GPR124. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except GPR125 (82 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Marmoset, Dog, Bat, Elephant, Panda, Horse, Rabbit, Pig, Opossum (100%) Mouse, Rat, Hamster, Turkey, Chicken, Xenopus (94%) Monkey, Bovine, Platypus, Lizard (82%).

Product Details

Purification: Immunoaffinity purified

Target Details

Target: GPR124

Alternative Name: ADGRA2 / GPR124 ([GPR124 Products](#))

Background: Name/Gene ID: ADGRA2

Subfamily: Orphan-U

Family: GPCR

Synonyms: ADGRA2, G protein-coupled receptor 124, G-protein coupled receptor 124, TEM5, GPR124, KIAA1531, Tumor endothelial marker 5

Gene ID: 25960

Application Details

Application Notes: Approved: IHC, IHC-P (4 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: Lot specific


Buffer: PBS, less than 0.1 % sodium azide.

Preservative: Sodium azide

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-GPR124 antibody ABIN1048704 IHC staining of human colon carcinoma, neoplastic cells. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-GPR124 / TEM5 antibody IHC of human Prostate, Carcinoma Endothelial Cells. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.