

Datasheet for ABIN1048719  
**anti-GPR137B antibody (C-Term)**



[Go to Product page](#)

3 Images

## Overview

Quantity:	50 µg
Target:	GPR137B
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Horse, Monkey, Pig, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPR137B antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 16 amino acid peptide from C-terminus of human GPR137B / TM7SF1. Percent identity with other species by BLAST analysis: Human, Gibbon, Monkey, Marmoset, Mouse, Rat, Panda, Horse, Rabbit, Pig, Opossum (100%), Hamster, Elephant, Bovine (94%), Goat, Chicken, Lizard (88%), Dog, Bat, Turkey (81%).  Type of Immunogen: Synthetic peptide
Specificity:	Human GPR137B / TM7SF1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gibbon, Monkey, Marmoset,

## Product Details

---

Mouse, Rat, Panda, Horse, Rabbit, Pig, Opossum (100%) Hamster, Elephant, Bovine (94%) Goat, Chicken, Lizard (88%) Dog, Bat, Turkey (81%).

Purification: Immunoaffinity purified

## Target Details

---

Target: GPR137B

Alternative Name: GPR137B ([GPR137B Products](#))

Background: Name/Gene ID: GPR137B  
Subfamily: Orphan-U  
Family: GPCR

Synonyms: GPR137B, TM7SF1

Gene ID: 7107

## Application Details

---

Application Notes: Approved: ICC, IHC, IHC-P (5 - 12 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### **Tissue Preparation**

Formalin fixation and embedding in paraffin wax

### **Tissue Sectioning**

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### **Rehydration**

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

### Handling

---

Format: Liquid

Concentration: Lot specific

Buffer: PBS, less than 0.1 % sodium azide.


Preservative: Sodium azide

---

## Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images




### Immunohistochemistry

**Image 1.** Anti-GPR137B antibody IHC staining of brain, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048719 dilution 41613 ug/ml.



### Immunohistochemistry

**Image 2.** Anti-GPR137B antibody IHC staining of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048719 dilution 41613 ug/ml.



### Immunocytochemistry

**Image 3.** Anti-GPR137B / TM7SF1 antibody immunocytochemistry (ICC) staining of HEK293 human embryonic kidney cells transfected with GPR137B / TM7SF1.