

Datasheet for ABIN1048789  
**anti-GPR3 antibody (Extracellular Domain)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	50 µg
Target:	GPR3
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Pig
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 15 amino acid peptide from 1st extracellular domain of human GPR3. Percent identity with other species by BLAST analysis: Human, Gorilla, Marmoset, Panda, Pig (100%), Gibbon, Monkey, Bovine, Elephant, Horse (93%), Mouse, Rat, Rabbit (87%).  Type of Immunogen: Synthetic peptide
Specificity:	Human GPR3. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except SPPL2A (60 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Marmoset, Panda, Pig (100%) Gibbon, Monkey, Bovine, Elephant, Horse (93%) Mouse, Rat, Rabbit (87%).
Purification:	Immunoaffinity purified

## Target Details

Target:	GPR3
Alternative Name:	GPR3 ( <a href="#">GPR3 Products</a> )
Background:	Name/Gene ID: GPR3 Subfamily: Lysophospholipid/Lysosphingolipid Family: GPCR  Synonyms: GPR3, ACCA, ACCA orphan receptor, G protein-coupled receptor 3, GPCR3, Hacca, G-protein coupled receptor 3
Gene ID:	2827
Pathways:	<a href="#">cAMP Metabolic Process</a>

## Application Details

Application Notes:	Approved: IHC, IHC-P (6 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	<b>The IHC-pro Immunohistochemistry Protocol</b>  <b>Tissue Preparation</b> Formalin fixation and embedding in paraffin wax  <b>Tissue Sectioning</b> Make 4-µm sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-drying oven for 45 minutes at 60°C Deparaffinization Wash slides in 3 changes of xylene – 5 minutes each at room temperature.  <b>Rehydration</b> Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature. Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature. Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature. Rinse slides in gentle running distilled water – 5 minutes at room temperature.  <b>Antigen retrieval</b> Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions:	For Research Use only
---------------	-----------------------

### Handling

---

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------

Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

Preservative:	Sodium azide
---------------	--------------

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
--------------------	--

---


## Handling

Storage: 4 °C, -20 °C

Storage Comment: Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.


Expiry Date: 12 months

## Images



### Immunohistochemistry

**Image 1.** Anti-GPR3 antibody ABIN1048789 IHC staining of human brain. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-GPR3 antibody IHC of human Ovary, Carcinoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.