



[Go to Product page](#)

Datasheet for ABIN1048868

## anti-GPR75 antibody (C-Term)

### 3 Images

#### Overview

Quantity:	50 µg
Target:	GPR75
Binding Specificity:	C-Term
Reactivity:	Human, Horse, Cow, Pig, Rabbit, Monkey, Bat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 19 amino acid peptide from C-terminus of human GPR75. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Bovine, Bat, Elephant, Horse, Rabbit, Pig (100%), Mouse, Rat, Hamster, Panda, Turkey (95%), Opossum, Chicken, Lizard, Xenopus (89%), Zebrafish (84%).  Type of Immunogen: Synthetic peptide
Specificity:	Human GPR75. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Bovine, Bat, Elephant, Horse, Rabbit, Pig (100%) Mouse, Rat, Hamster, Panda, Turkey (95%) Opossum, Chicken, Lizard, Xenopus (89%) Zebrafish (84%).

## Product Details

---

Purification: Immunoaffinity purified

## Target Details

---

Target: GPR75

Alternative Name: GPR75 ([GPR75 Products](#))

Background: Name/Gene ID: GPR75  
Subfamily: Orphan-A  
Family: GPCR

Synonyms: GPR75, G protein-coupled receptor 75, GPR-chr2, WI31133, GPRchr2, WI-31133

Gene ID: 10936

## Application Details

---

Application Notes: Approved: IHC, IHC-P (10 - 25 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### **Tissue Preparation**

Formalin fixation and embedding in paraffin wax

### **Tissue Sectioning**

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### **Rehydration**

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### **Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### **Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### **Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: Lot specific

Buffer: PBS, less than 0.1 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

---

## Handling


should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.


Expiry Date: 12 months

## Images




### Immunohistochemistry

**Image 1.** Anti-GPR75 antibody IHC staining of human brain, cortex, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048868 dilution 45931 ug/ml.



### Immunohistochemistry

**Image 2.** Anti-GPR75 antibody IHC staining of human testis. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1048868 dilution 45931 ug/ml.



### Immunohistochemistry

**Image 3.** Anti-GPR75 antibody IHC of human brain, cortex, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 10-25 ug/ml.