

Datasheet for ABIN1048896

**anti-GPR89A antibody (Extracellular Domain)****2** Images[Go to Product page](#)

## Overview

Quantity:	50 µg
Target:	GPR89A
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Mouse, Rat, Cow, Horse, Dog, Zebrafish (Danio rerio), Pig, Hamster, Monkey, Bat, Chicken, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPR89A antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC)

## Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 16 amino acid peptide from 3rd extracellular domain of human GPR89A. Percent identity with other species by BLAST analysis: Human, Gibbon, Monkey, Marmoset, Mouse, Rat, Dog, Bat, Bovine, Hamster, Panda, Horse, Pig, Opossum, Chicken, Platypus, Xenopus, Salmon, Pufferfish, Zebrafish, Stickleback, Sea anemone (100%), Capsaspora (94%), Eye worm, Mosquito, Ant, Beetle, Trichina worm, Trichoplax (88%), Nematode (81%).  Type of Immunogen: Synthetic peptide
Specificity:	Human GPR89A. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

## Product Details

Predicted Reactivity: Percent identity with other species by BLAST analysis: Human, Gibbon, Monkey, Marmoset, Mouse, Rat, Dog, Bat, Bovine, Hamster, Panda, Horse, Pig, Opossum, Chicken, Platypus, Xenopus, Salmon, Pufferfish, Zebrafish, Stickleback, Sea anemone (100%) Capsaspora (94%) Eye worm, Mosquito, Ant, Beetle, Trichina worm, Trichoplax (88%) Nematode (81%).

Purification: Immunoaffinity purified

## Target Details

Target: GPR89A

Alternative Name: GPR89A ([GPR89A Products](#))

Background: Name/Gene ID: GPR89A  
Subfamily: Orphan-A  
Family: GPCR

Synonyms: GPR89A, G protein-coupled receptor 89A, Golgi pH regulator A, GPHRA, GPR89, Protein GPR89, Protein GPR89A, SH120, UNQ192

Gene ID: 653519

## Application Details

Application Notes: Approved: IHC, IHC-P (10 - 20 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions:	For Research Use only
---------------	-----------------------

### Handling

---

Format:	Liquid
---------	--------


Concentration:	Lot specific
----------------	--------------

---

## Handling


Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images



### Immunohistochemistry

**Image 1.** Anti-GPR89A antibody ABIN1048896 IHC staining of human brain, hippocampus. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunohistochemistry

**Image 2.** Anti-GPR89A antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.