antibodies -online.com





anti-GPRC5A antibody (Transmembrane Domain)

2 Images



Go to Product page

_							
0	V	е	r١	/1	е	V	I

Quantity:	50 μg
Target:	GPRC5A
Binding Specificity:	Transmembrane Domain
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Brand:	IHC-plus™
Immunogen:	Synthetic 15 amino acid peptide from 3rd transmembrane domain of human GPRC5A / RAI3. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset (100%), Panda, Dog, Rabbit (93%), Mouse, Hamster, Bovine (87%), Elephant, Horse (80%).
	Type of Immunogen: Synthetic peptide
Specificity:	
,	Human GPRC5A / RAI3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Product Details Purification: Immunoaffinity purified Target Details Target: GPRC5A Alternative Name: GPRC5A / RAI3 (GPRC5A Products)

Name/Gene ID: GPRC5A

Subfamily: Orphan-C

Family: GPCR

Synonyms: GPRC5A, RAIG-1, Retinoic acid induced 3, Retinoic acid responsive, GPCR5A, RAI3,

RAIG1, Retinoic acid-inducible gene 1

Gene ID: 9052

Application Details

Background:

Application Notes:	Approved: IHC, IHC-P (16 - 17 μg/mL)
Comment:	Target Species of Antibody: Human

Assay Procedure:

The IHC-pro Immunohistochemistry Protocol

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene - 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol - 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol - 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only


Handling

Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, less than 0.1 % sodium azide.	
Preservative:	Sodium azide	

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-GPRC5A / RAI3 antibody ABIN1048909 IHC staining of human breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-GPRC5A / RAI3 antibody IHC of human testis. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval.