

Datasheet for ABIN1048934

## anti-Metabotropic Glutamate Receptor 3 antibody (N-Term)



[Go to Product page](#)

### 3 Images

#### Overview

Quantity:	50 µg
Target:	Metabotropic Glutamate Receptor 3 (GRM3)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Horse, Monkey, Hamster, Pig
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 16 amino acid peptide from N-terminal extracellular domain of human GRM3 / MGLUR3. Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Horse, Pig, Opossum, Platypus, Lizard (100%), Rabbit, Turkey, Zebra finch, Chicken (94%).  Type of Immunogen: Synthetic peptide
Specificity:	Human GRM3 / MGLUR3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Orangutan, Gibbon, Monkey, Marmoset, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Horse, Pig, Opossum, Platypus, Lizard (100%) Rabbit, Turkey, Zebra finch, Chicken (94%).

## Product Details

---

Purification: Immunoaffinity purified

## Target Details

---

Target: Metabotropic Glutamate Receptor 3 (GRM3)

Alternative Name: GRM3 / MGLUR3 ([GRM3 Products](#))

Background: Name/Gene ID: GRM3  
Subfamily: Metabotropic glutamate  
Family: GPCR

Synonyms: GRM3, GPRC1C, GLUR3, MGLUR3, MGlu3

Gene ID: 2913

Pathways: [cAMP Metabolic Process](#), [Synaptic Membrane](#)

## Application Details

---

Application Notes: Approved: ICC, IHC, IHC-P (16 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### **Tissue Preparation**

Formalin fixation and embedding in paraffin wax

### **Tissue Sectioning**

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### **Rehydration**

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions: For Research Use only

### Handling

---

Format: Liquid

Concentration: Lot specific

Buffer: PBS, less than 0.1 % sodium azide.


Preservative: Sodium azide

---

## Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images




### Immunohistochemistry

**Image 1.** Anti-GRM3 / MGLUR3 antibody ABIN1048934 IHC staining of human brain, pituitary. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



### Immunocytochemistry

**Image 2.** Anti-GRM3 / MGLUR3 antibody immunocytochemistry (ICC) staining of untransfected HEK293 human embryonic kidney cells.



### Immunohistochemistry

**Image 3.** Anti-GRM3 / MGLUR3 antibody IHC of human Skin, Melanoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.