

Datasheet for ABIN1048943
anti-GRM7 antibody (N-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	50 µg
Target:	GRM7
Binding Specificity:	N-Term
Reactivity:	Human, Cow
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 19 amino acid peptide from N-terminal extracellular domain of human GRM7 / MGLUR7. Percent identity with other species by BLAST analysis: Human, Elephant, Bovine (100%), Chimpanzee, Gorilla, Orangutan, Monkey, Marmoset, Mouse, Rat, Dog, Horse (95%), Gibbon, Panda, Rabbit, Opossum (89%), Chicken, Xenopus (84%). Type of Immunogen: Synthetic peptide
Specificity:	Human GRM7 / MGLUR7. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except GRM8 (63 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Elephant, Bovine (100%) Chimpanzee, Gorilla, Orangutan, Monkey, Marmoset, Mouse, Rat, Dog, Horse (95%) Gibbon, Panda, Rabbit, Opossum (89%) Chicken, Xenopus (84%).

Product Details

Purification: Immunoaffinity purified

Target Details

Target: GRM7

Alternative Name: GRM7 / MGLUR7 ([GRM7 Products](#))

Background: Name/Gene ID: GRM7
Subfamily: Metabotropic glutamate
Family: GPCR

Synonyms: GRM7, GLUR7, MGLU7, MGLUR7, GPRC1G

Gene ID: 2917

Pathways: [Sensory Perception of Sound](#), [cAMP Metabolic Process](#), [Feeding Behaviour](#)

Application Details

Application Notes: Approved: ICC, IHC, IHC-P (6 - 16 µg/mL), WB

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------


Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

Preservative:	Sodium azide
---------------	--------------

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-GRM7 / MGLUR7 antibody ABIN1048943 IHC staining of human brain, neurons and glia. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-GRM7 / MGLUR7 antibody IHC of human adrenal. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.