.-online.com antibodies

Datasheet for ABIN1048956 anti-HRH1 antibody (Extracellular Domain)





Overview

Quantity:	50 µg
Target:	HRH1
Binding Specificity:	Extracellular Domain
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 17 amino acid peptide from 2nd extracellular domain of human HRH1 / Histamine H1 Receptor. Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey (100%), Marmoset (94%), Orangutan (88%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human HRH1 / Histamine H1 Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey (100%) Marmoset (94%) Orangutan (88%).
Purification:	Immunoaffinity purified

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN1048956 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	HRH1
Alternative Name:	HRH1 / Histamine H1 Receptor (HRH1 Products)
Background:	Name/Gene ID: HRH1
	Subfamily: Histamine
	Family: GPCR
	Synonyms: HRH1, H1 histamine receptor, H1 receptor, Histamine 1 receptor, Histamine H1
	receptor, H1-R, H1R, Histamine H(1) receptor, Histamine receptor H1, HisH1
Gene ID:	3269
Pathways:	Regulation of Carbohydrate Metabolic Process
Application Details	
Application Notes:	Approved: IHC, IHC-P (11 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	The IHC-pro Immunohistochemistry Protocol
	Tissue Preparation
	Formalin fixation and embedding in paraffin wax
	Tissue Sectioning
	Make 4-µm sections and place on pre-cleaned and charged microscope slides.
	Heat in a tissue-drying oven for 45 minutes at 60°C
	Deparaffinization
	Wash slides in 3 changes of xylene – 5 minutes each at room temperature.
	Rehydration
	Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.
	Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.
	Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.
	Rinse slides in gentle running distilled water – 5 minutes at room temperature.
	Antigen retrieval
	Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN1048956 | 09/12/2023 | Copyright antibodies-online. All rights reserved. Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure. Apply a universal protein block – 20 minutes at room temperature. Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature. Rinse slides in 1X TBST - 1 minute at room temperature. Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature. Rinse slides 1X TBST – 1 minute at room temperature. Apply alkaline phosphatase streptavidin – 30 minutes at room temperature. Rinse slides in 1X TBST - 1 minute at room temperature. Apply alkaline phosphatase streptavidin – 30 minutes at room temperature. May alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble. Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature. Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature. Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature. Wash slides in 3 changes of xylene – 1 minute each at room temperature. Apply coverslip

Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN1048956 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Handling


Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-HRH1 / Histamine H1 Receptor antibody ABIN1048956 IHC staining of human brain, neurons and glia. Immunohistochemistry of formalin-fixed, paraffinembedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-HRH1 / Histamine H1 Receptor antibody IHC of human Vessel. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN1048956 | 09/12/2023 | Copyright antibodies-online. All rights reserved.