.-online.com antibodies

Datasheet for ABIN1048962 anti-HRH3 antibody (Cytoplasmic Domain)

5 Images



Overview

Quantity:	50 µg
Target:	HRH3
Binding Specificity:	Cytoplasmic Domain
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 16 amino acid peptide from 3rd cytoplasmic domain of human HRH3 / Histamine H3 Receptor. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey (100%), Mouse, Rat, Hamster, Elephant, Bovine, Dog, Horse, Opossum, Guinea pig, Platypus (88%), Bat, Turkey (81%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human HRH3 / Histamine H3 Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey (100%) Mouse, Rat, Hamster, Elephant, Bovine, Dog, Horse, Opossum, Guinea pig, Platypus (88%) Bat, Turkey (81%).

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/5 | Product datasheet for ABIN1048962 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Product Details

Purification:

Immunoaffinity purified

Target Details

Target:	HRH3
Alternative Name:	HRH3 / Histamine 3 Receptor (HRH3 Products)
Background:	Name/Gene ID: HRH3 Subfamily: Histamine Family: GPCR
	Synonyms: HRH3, G protein-coupled receptor 97, GPCR97, H3 histamine receptor, H3R, HH3R, Histamine H3 receptor, Histamine receptor H3, G-protein coupled receptor 97, Histamine 3 receptor
Gene ID:	11255
Pathways:	cAMP Metabolic Process, Feeding Behaviour

Application Details

Application Notes:	Approved: ICC, IHC, IHC-P (18 - 36 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	The IHC-pro Immunohistochemistry Protocol
	Tissue Preparation Formalin fixation and embedding in paraffin wax
	Tissue Sectioning Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene - 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature. Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature. Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/5 | Product datasheet for ABIN1048962 | 09/12/2023 | Copyright antibodies-online. All rights reserved. Rinse slides in gentle running distilled water - 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure. Apply a universal protein block – 20 minutes at room temperature. Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature. Rinse slides in 1X TBST - 1 minute at room temperature. Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature. Rinse slides 1X TBST – 1 minute at room temperature. Apply alkaline phosphatase streptavidin – 30 minutes at room temperature. Rinse slides in 1X TBST - 1 minute at room temperature. Apply alkaline phosphatase streptavidin – 30 minutes at room temperature. Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble. Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature. Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature. Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature. Wash slides in 3 changes of xylene – 1 minute each at room temperature. Apply coverslip

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/5 | Product datasheet for ABIN1048962 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

Handling

Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images




Image 1.




Image 2.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/5 | Product datasheet for ABIN1048962 | 09/12/2023 | Copyright antibodies-online. All rights reserved.




Image 3.

Please check the product details page for more images. Overall 5 images are available for ABIN1048962.