

Datasheet for ABIN1048968

**anti-Serotonin Receptor 6 antibody (C-Term)**

3 Images

[Go to Product page](#)

## Overview

Quantity:	50 µg
Target:	Serotonin Receptor 6 (HTR6)
Binding Specificity:	C-Term
Reactivity:	Human, Rat, Monkey, Dog, Hamster, Horse
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

## Product Details

Brand:	IHC-plus™
Immunogen:	<p>Synthetic 17 amino acid peptide from C-terminus of human HTR6 / 5-HT36. Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey, Marmoset, Rat, Hamster, Panda, Dog, Horse (100%), Mouse, Elephant, Bovine, Bat, Rabbit (94%), Stickleback, Pufferfish (88%), Turkey, Chicken, Xenopus, Zebrafish (82%).</p> <p>Type of Immunogen: Synthetic peptide</p>
Specificity:	Human HTR6 / 5-HT36. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey, Marmoset, Rat, Hamster, Panda, Dog, Horse (100%) Mouse, Elephant, Bovine, Bat, Rabbit (94%) Stickleback, Pufferfish (88%) Turkey, Chicken, Xenopus, Zebrafish (82%).

## Product Details

Purification: Immunoaffinity purified

## Target Details

Target: Serotonin Receptor 6 (HTR6)

Alternative Name: HTR6 / 5-HT6 Receptor ([HTR6 Products](#))

Background: Name/Gene ID: HTR6

Subfamily: Serotonin

Family: GPCR

Synonyms: HTR6, 5-HT6, 5-HT6R, 5-HT-6, 5-hydroxytryptamine receptor 6, 5-HT6 receptor, 5-HT6 serotonin receptor, 5-hydroxytryptamine 6 receptor, Serotonin 6 receptor, Serotonin receptor 6, Serotonin 5-HT-6 receptor, Serotonin type 6 receptor

Gene ID: 3362

Pathways: [JAK-STAT Signaling](#)

## Application Details

Application Notes: Approved: IHC, IHC-P (10 - 20 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

### Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

### Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

### Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

---

Restrictions:	For Research Use only
---------------	-----------------------

### Handling

---

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------


Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

---

## Handling


Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images




### Immunohistochemistry

**Image 1.** Human, Brain, Cerebellum: Formalin-Fixed Paraffin-Embedded (FFPE)



### Immunohistochemistry

**Image 2.** Anti-HTR6 / 5-HT6 antibody ABIN1048968 IHC staining of human brain, neuron. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



#### Immunohistochemistry

**Image 3.** Human, Brain, Cerebellum: Formalin-Fixed Paraffin-Embedded (FFPE)