

Datasheet for ABIN1049068

anti-MRGPRD antibody (Extracellular Domain)

2 Images



Go to Product page

\sim		
()\//	erv	$\square \square \square$
\cup	_I V	$I \subset V \setminus$

Quantity:	50 μg
Target:	MRGPRD
Binding Specificity:	Extracellular Domain
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MRGPRD antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Brand:	IHC-plus™
lmmunogen:	Synthetic 18 amino acid peptide from 2nd extracellular domain of human MRGPRD. Percent identity with other species by BLAST analysis: Human (100%), Gorilla, Gibbon, Monkey (94%).
	Type of Immunogen: Synthetic peptide
Specificity:	Human MRGPRD. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human (100%) Gorilla, Gibbon, Monkey (94%).
Purification:	Immunoaffinity purified

Target Details

Target:	MRGPRD
Alternative Name:	MRGD / MRGPRD (MRGPRD Products)
Background:	Name/Gene ID: MRGPRD
	Subfamily: Orphan-A
	Family: GPCR
	Synonyms: MRGPRD, Beta-alanine receptor, MRGD, MAS-related GPR, member D, TGR7
Gene ID:	116512

Application Details

Assay Procedure:	The IHC-pro Immunohistochemistry Protocol
Comment:	Target Species of Antibody: Human
Application Notes:	Approved: IHC, IHC-P (9.3 μg/mL)

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol - 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol - 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for

up to 1 month. Avoid freeze-thaw cycles.

Expiry Date:


12 months

Images



Immunohistochemistry

Image 1. Anti-MRGPRD antibody ABIN1049068 IHC staining of human brain, medulla. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-MRGD / MRGPRD antibody IHC of human Skin, Melanoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.