

Datasheet for ABIN1049083
anti-MST1R antibody (N-Term)[Go to Product page](#)

1 Image

Overview

Quantity:	50 µg
Target:	MST1R
Binding Specificity:	N-Term
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 18 amino acid peptide from N-terminus of human MST1R / RON. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset (100%), Gibbon, Dog (94%), Rabbit (89%), Panda, Bovine (83%). Type of Immunogen: Synthetic peptide
Specificity:	Human MST1R / RON. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset (100%) Gibbon, Dog (94%) Rabbit (89%) Panda, Bovine (83%).
Purification:	Immunoaffinity purified

Target Details

Target:	MST1R
Alternative Name:	MST1R / RON (MST1R Products)
Background:	<p>Name/Gene ID: MST1R</p> <p>Subfamily: HGF Receptor/MET</p> <p>Family: Protein Kinase</p> <p>Synonyms: MST1R, C-met-related tyrosine kinase, CD136, CDw136, CD136 antigen, Friend virus susceptibility 2, Fv2, MSP receptor, p185-Ron, PTK8 protein tyrosine kinase 8, RON, Soluble RON variant 3, Soluble RON variant 1, MST1R variant RON30, Oncogene RON, PTK8, MST1R variant RON62, Protein-tyrosine kinase 8, RON variant E2E3, Soluble RON variant 2, Soluble RON variant 4</p>
Gene ID:	4486
Pathways:	RTK Signaling

Application Details

Application Notes:	Approved: IHC, IHC-P (2.5 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	<p>The IHC-pro Immunohistochemistry Protocol</p> <p>Tissue Preparation</p> <p>Formalin fixation and embedding in paraffin wax</p> <p>Tissue Sectioning</p> <p>Make 4-µm sections and place on pre-cleaned and charged microscope slides.</p> <p>Heat in a tissue-drying oven for 45 minutes at 60°C</p> <p>Deparaffinization</p> <p>Wash slides in 3 changes of xylene – 5 minutes each at room temperature.</p> <p>Rehydration</p> <p>Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.</p> <p>Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.</p> <p>Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.</p> <p>Rinse slides in gentle running distilled water – 5 minutes at room temperature.</p>

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------


Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

Preservative:	Sodium azide
---------------	--------------

Handling

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-MST1R / RON antibody ABIN1049083 IHC staining of human colon, surface epithelium. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.