

Datasheet for ABIN1049128

anti-NR0B2 antibody (Ligand Binding Domain)[Go to Product page](#)**2** Images

Overview

| | |
|----------------------|--|
| Quantity: | 50 µg |
| Target: | NR0B2 |
| Binding Specificity: | Ligand Binding Domain |
| Reactivity: | Human, Mouse, Rabbit, Monkey, Hamster |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Application: | Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|-----------------------|---|
| Brand: | IHC-plus™ |
| Immunogen: | Synthetic 16 amino acid peptide from ligand-binding domain of human NR0B2. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Mouse, Hamster, Rabbit (100%), Rat, Bovine, Bat (94%), Elephant, Panda, Horse, Pig, Opossum (88%), Dog (81%). Type of Immunogen: Synthetic peptide |
| Specificity: | Human NR0B2. BLAST analysis of the peptide immunogen showed no homology with other human proteins. |
| Predicted Reactivity: | Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Mouse, Hamster, Rabbit (100%) Rat, Bovine, Bat (94%) Elephant, Panda, Horse, Pig, Opossum (88%) Dog (81%). |

Product Details

Purification: Immunoaffinity purified

Target Details

Target: NR0B2

Alternative Name: NR0B2 / SHP ([NR0B2 Products](#))

Background: Name/Gene ID: NR0B2
Subfamily: NR0 Knirps-like
Family: NHR

Synonyms: NR0B2, Short heterodimer partner, SHP, Nuclear receptor SHP, Orphan nuclear receptor SHP, Small heterodimer partner, SHP1

Gene ID: 8431

Pathways: [Nuclear Receptor Transcription Pathway](#), [Positive Regulation of Peptide Hormone Secretion](#), [Intracellular Steroid Hormone Receptor Signaling Pathway](#), [Steroid Hormone Mediated Signaling Pathway](#)

Application Details

Application Notes: Approved: ICC, IF, IHC, IHC-P (4 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling


| | |
|---------|--------|
| Format: | Liquid |
|---------|--------|

| | |
|----------------|--------------|
| Concentration: | Lot specific |
|----------------|--------------|

Handling


| | |
|--------------------|---|
| Buffer: | PBS, less than 0.1 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles. |
| Expiry Date: | 12 months |

Images



Immunohistochemistry

Image 1. Anti-NR0B2 antibody ABIN1049128 IHC staining of human liver, hepatocytes. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-NR0B2 antibody IHC of human liver, hepatocytes. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.