

Datasheet for ABIN1049142
anti-NR4A3 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	50 µg
Target:	NR4A3
Binding Specificity:	C-Term
Reactivity:	Human, Dog, Bat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 20 amino acid peptide from near C-terminus of human NR4A3. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Dog, Bat, Elephant, Panda (100%), Marmoset, Rat, Bovine, Hamster, Rabbit, Pig (95%), Mouse (90%). Type of Immunogen: Synthetic peptide
Specificity:	Human NR4A3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Dog, Bat, Elephant, Panda (100%) Marmoset, Rat, Bovine, Hamster, Rabbit, Pig (95%) Mouse (90%).
Purification:	Immunoaffinity purified

Target Details

Target:	NR4A3
Alternative Name:	Nor-1 / NR4A3 (NR4A3 Products)
Background:	<p>Name/Gene ID: NR4A3</p> <p>Subfamily: NR4 Nerve growth factor-like</p> <p>Family: NHR</p> <p>Synonyms: NR4A3, CHN, MINOR, NOR1, Neuron-derived orphan receptor, Nuclear hormone receptor NOR-1, Nor-1, CSMF, Orphan receptor nor1, TEC receptor</p>
Gene ID:	8013
Pathways:	Fc-epsilon Receptor Signaling Pathway , Nuclear Receptor Transcription Pathway , Steroid Hormone Mediated Signaling Pathway

Application Details

Application Notes:	Approved: IHC, IHC-P (10 - 20 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	<p>The IHC-pro Immunohistochemistry Protocol</p> <p>Tissue Preparation</p> <p>Formalin fixation and embedding in paraffin wax</p> <p>Tissue Sectioning</p> <p>Make 4-µm sections and place on pre-cleaned and charged microscope slides.</p> <p>Heat in a tissue-drying oven for 45 minutes at 60°C</p> <p>Deparaffinization</p> <p>Wash slides in 3 changes of xylene – 5 minutes each at room temperature.</p> <p>Rehydration</p> <p>Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.</p> <p>Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.</p> <p>Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.</p> <p>Rinse slides in gentle running distilled water – 5 minutes at room temperature.</p> <p>Antigen retrieval</p>

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------

Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------


Preservative:	Sodium azide
---------------	--------------

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
--------------------	---

Handling


	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-NR4A3 antibody ABIN1049142 IHC staining of human brain, neurons and glia. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-Nor-1 / NR4A3 antibody IHC of human Skin, Melanoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.