

Datasheet for ABIN1049168

anti-OPN5 antibody (C-Term)





Go to Product page

	ve	rv	ie	W
\circ	v C	· I V	10	V V

Quantity:	50 μg
Target:	OPN5
Binding Specificity:	C-Term
Reactivity:	Human, Rabbit, Cow, Dog, Horse, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OPN5 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Brand:	IHC-plus™
Brand: Immunogen:	IHC-plus™ Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other
	Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other
	Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Dog, Bovine, Elephant, Horse,
	Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Dog, Bovine, Elephant, Horse, Rabbit (100%), Gibbon, Mouse (94%), Bat (81%).
Immunogen:	Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Dog, Bovine, Elephant, Horse, Rabbit (100%), Gibbon, Mouse (94%), Bat (81%). Type of Immunogen: Synthetic peptide
Immunogen:	Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Dog, Bovine, Elephant, Horse, Rabbit (100%), Gibbon, Mouse (94%), Bat (81%). Type of Immunogen: Synthetic peptide Human OPN5. BLAST analysis of the peptide immunogen showed no homology with other
Immunogen: Specificity:	Synthetic 16 amino acid peptide from C-terminus of human OPN5. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey, Marmoset, Dog, Bovine, Elephant, Horse, Rabbit (100%), Gibbon, Mouse (94%), Bat (81%). Type of Immunogen: Synthetic peptide Human OPN5. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Target Details

Target:	OPN5
Alternative Name:	OPN5 (OPN5 Products)
Background:	Name/Gene ID: OPN5
	Subfamily: Orphan-A
	Family: GPCR
	Synonyms: OPN5, G-protein coupled receptor 136, Neuropsin, Opsin 5, Opsin-5, PGR12,
	TMEM13, Transmembrane protein 13, DJ402H5.1, GPR136
Gene ID:	221391

Application Details

Assay Procedure:	The IHC-pro Immunohistochemistry Protocol
Comment:	Target Species of Antibody: Human
Application Notes:	Approved: IHC, IHC-P (16 - 32 μg/mL)

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol - 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate - 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol - 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene - 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C


Handling

Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for

up to 1 month. Avoid freeze-thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Anti-OPN5 antibody ABIN1049168 IHC staining of human testis. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-OPN5 antibody IHC of human Ovary, Carcinoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.