

Datasheet for ABIN1049188

## anti-OR6N1 antibody (Extracellular Domain)



[Go to Product page](#)

### 3 Images

#### Overview

Quantity:	50 µg
Target:	OR6N1
Binding Specificity:	Extracellular Domain
Reactivity:	Human, Rat, Mouse, Hamster, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR6N1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

#### Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 15 amino acid peptide from 1st extracellular domain of human OR6N1. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Rabbit, Opossum (100%), Marmoset, Dog (93%), Panda, Bovine, Horse, Pig (87%).  Type of Immunogen: Synthetic peptide
Specificity:	Human OR6N1. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except OR6K3 (67 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Rabbit, Opossum (100%) Marmoset, Dog (93%) Panda, Bovine, Horse,

## Product Details

Pig (87%).

Purification: Immunoaffinity purified

## Target Details

Target: OR6N1

Alternative Name: OR6N1 ([OR6N1 Products](#))

Background: Name/Gene ID: OR6N1  
Subfamily: Olfactory  
Family: GPCR

Synonyms: OR6N1, Olfactory receptor OR1-22, Olfactory receptor 6N1, OR1-22

Gene ID: 128372

## Application Details

Application Notes: Approved: IHC, IHC-P (7 µg/mL)

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

### Tissue Preparation

Formalin fixation and embedding in paraffin wax

### Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

### Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

**Antigen retrieval**

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes  
Remove from heat and let stand at room temperature in buffer - 20 minutes  
Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

**Immunostaining**

Do not allow tissues to dry at any time during the staining procedure.  
Apply a universal protein block – 20 minutes at room temperature.  
Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.  
Rinse slides in 1X TBST - 1 minute at room temperature.  
Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.  
Rinse slides 1X TBST – 1 minute at room temperature.  
Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.  
Rinse slides in 1X TBST - 1 minute at room temperature.  
Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.  
Wash slides in distilled water – 1 minute at room temperature.

**Dehydrate**

This method should only be used if the chromogen substrate is alcohol insoluble.  
Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.  
Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.  
Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.  
Wash slides in 3 changes of xylene – 1 minute each at room temperature.  
Apply coverslip

---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

---


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide

---

## Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

## Images




### Immunohistochemistry

**Image 1.** Anti-OR6N1 antibody IHC staining of human brain, cortex, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1049188 concentration 41791 ug/ml.



### Immunohistochemistry

**Image 2.** Anti-OR6N1 antibody IHC staining of human brain, cerebellum, Purkinje neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody ABIN1049188 concentration 41791 ug/ml.



#### Immunohistochemistry

**Image 3.** Anti-OR6N1 antibody IHC of human brain, cortex, neurons. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 6-14 ug/ml.