

Datasheet for ABIN1049281 anti-PYGM antibody (Internal Region)



[Go to Product page](#)

2 Images

Overview

Quantity:	50 µg
Target:	PYGM
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat, Rabbit, Cow, Dog, Sheep, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PYGM antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	<p>Synthetic 10 amino acid peptide from internal region of human PYGM / Phosphorylase b.</p> <p>Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Baboon, Monkey, Galago, Marmoset, Mouse, Rat, Sheep, Dog, Bovine, Elephant, Panda, Rabbit, Armadillo (100%), Shrew, Bat, Pig, Platypus (90%), Xenopus, Penicillium (80%).</p> <p>Type of Immunogen: Synthetic phospho-peptide</p>
Specificity:	Human PYGM / Phosphorylase b. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Baboon, Monkey,

Product Details

Galago, Marmoset, Mouse, Rat, Sheep, Dog, Bovine, Elephant, Panda, Rabbit, Armadillo (100%)
Shrew, Bat, Pig, Platypus (90%) Xenopus, Penicillium (80%).

Purification: Immunoaffinity purified

Target Details

Target: PYGM

Alternative Name: PYGM ([PYGM Products](#))

Background: Name/Gene ID: PYGM

Synonyms: PYGM, GPMM, Myophosphorylase, Phosphorylase, glycogen muscle

Gene ID: 5837

Pathways: [Cellular Glucan Metabolic Process](#)

Application Details

Application Notes: Approved: IHC, IHC-P (10 µg/mL), WB

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes
Remove from heat and let stand at room temperature in buffer - 20 minutes
Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.
Apply a universal protein block – 20 minutes at room temperature.
Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.
Rinse slides in 1X TBST - 1 minute at room temperature.
Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.
Rinse slides 1X TBST – 1 minute at room temperature.
Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.
Rinse slides in 1X TBST - 1 minute at room temperature.
Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.
Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.
Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.
Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.
Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.
Wash slides in 3 changes of xylene – 1 minute each at room temperature.
Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-PYGM / Phosphorylase b antibody ABIN1049281 IHC staining of human skeletal muscle. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-PYGM / Phosphorylase b antibody IHC of human skeletal muscle. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.