

Datasheet for ABIN1049311

anti-Relaxin 3 Receptor 1 antibody (Cytoplasmic Domain)[Go to Product page](#)**2** Images

Overview

Quantity:	50 µg
Target:	Relaxin 3 Receptor 1 (RXFP3)
Binding Specificity:	Cytoplasmic Domain
Reactivity:	Human, Rat, Hamster
Host:	Rabbit
Clonality:	Polyclonal
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 20 amino acid peptide from 2nd cytoplasmic domain of human RXFP3. Percent identity with other species by BLAST analysis: Human, Gorilla, Marmoset, Rat, Hamster (100%), Monkey, Mouse, Bovine, Panda, Pig (95%), Rabbit, Opossum (80%). Type of Immunogen: Synthetic peptide
Specificity:	Human RXFP3. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except VGLL4 (45 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Marmoset, Rat, Hamster (100%) Monkey, Mouse, Bovine, Panda, Pig (95%) Rabbit, Opossum (80%).
Purification:	Immunoaffinity purified

Target Details

Target:	Relaxin 3 Receptor 1 (RXFP3)
Alternative Name:	RXFP3 (RXFP3 Products)
Background:	Name/Gene ID: RXFP3 Subfamily: Relaxin Family: GPCR Synonyms: RXFP3, GPCR135, Relaxin-3 receptor 1, Relaxin 3 receptor 1, RXFPR3, RLN3R1, SALPR, RLN3 receptor 1
Gene ID:	51289

Application Details

Application Notes:	Approved: IHC, IHC-P (14 - 28 µg/mL)
Comment:	Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	Lot specific
----------------	--------------

Buffer:	PBS, less than 0.1 % sodium azide.
---------	------------------------------------

Preservative:	Sodium azide
---------------	--------------

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
--------------------	--


Storage:	4 °C,-20 °C
----------	-------------

Handling

Storage Comment: Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Anti-RXFP3 antibody ABIN1049311 IHC staining of human brain, neurons and glia. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-RXFP3 antibody IHC of human Skin, Melanoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.