

Datasheet for ABIN1049373
anti-SYNGR2 antibody (Internal Region)



[Go to Product page](#)

3 Images

Overview

Quantity:	50 µg
Target:	SYNGR2
Binding Specificity:	Internal Region
Reactivity:	Human, Rat, Mouse, Cow, Hamster, Horse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SYNGR2 antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (IHC)

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 18 amino acid peptide from internal region of human SYNGR2 / Synaptogyrin 2. Percent identity with other species by BLAST analysis: Human, Orangutan, Gibbon, Monkey, Mouse, Rat, Hamster, Panda, Bovine, Horse (100%), Marmoset, Elephant, Guinea pig (94%), Galago, Dog, Rabbit (89%), Opossum (83%). Type of Immunogen: Synthetic peptide
Specificity:	Human SYNGR2 / Synaptogyrin 2. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Orangutan, Gibbon, Monkey, Mouse, Rat, Hamster, Panda, Bovine, Horse (100%) Marmoset, Elephant, Guinea pig (94%)

Product Details

Galago, Dog, Rabbit (89%) Opossum (83%).

Purification: Immunoaffinity purified

Target Details

Target: SYNGR2

Alternative Name: SYNGR2 / Synaptogyrin 2 ([SYNGR2 Products](#))

Background: Name/Gene ID: SYNGR2

Synonyms: SYNGR2, Cellugyrin, Synaptogyrin 2, Synaptogyrin-2

Gene ID: 9144

Application Details

Application Notes: Approved: IHC, IHC-P (15 - 20 µg/mL)

Usage: Immunohistochemistry with formalin-fixed paraffin-embedded tissues requires pretreatment with Proteinase K.

Comment: Target Species of Antibody: Human

Assay Procedure: **The IHC-pro Immunohistochemistry Protocol**

Tissue Preparation

Formalin fixation and embedding in paraffin wax

Tissue Sectioning

Make 4-µm sections and place on pre-cleaned and charged microscope slides.

Heat in a tissue-drying oven for 45 minutes at 60°C

Deparaffinization

Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: Lot specific


Buffer: PBS, less than 0.1 % sodium azide.

Preservative: Sodium azide

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images




Immunohistochemistry

Image 1. Anti-SYNGR2 / Synaptogyrin 2 antibody ABIN1049373 IHC staining of human, breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-SYNGR2 / Synaptogyrin 2 antibody ABIN1049373 IHC staining of human, prostate. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 3. Anti-SYNGR2 / Synaptogyrin 2 antibody IHC of human, breast. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.