

Datasheet for ABIN1049480 anti-WNT6 antibody (N-Term)

1 Image



Go to Product page

()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Quantity:	50 μg
Target:	WNT6
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog, Rabbit, Horse, Hamster
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WNT6 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Brand:	IHC-plus™
	πο ριασ
Immunogen:	Synthetic 20 amino acid peptide from N-terminus of human WNT6. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Rabbit (100%), Marmoset, Bat, Opossum, Platypus (95%).
Immunogen:	Synthetic 20 amino acid peptide from N-terminus of human WNT6. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant,
Immunogen: Specificity:	Synthetic 20 amino acid peptide from N-terminus of human WNT6. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Rabbit (100%), Marmoset, Bat, Opossum, Platypus (95%).
	Synthetic 20 amino acid peptide from N-terminus of human WNT6. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Mouse, Rat, Hamster, Elephant, Panda, Bovine, Dog, Horse, Rabbit (100%), Marmoset, Bat, Opossum, Platypus (95%). Type of Immunogen: Synthetic peptide Human WNT6. BLAST analysis of the peptide immunogen showed no homology with other

Product Details Purification: Immunoaffinity purified **Target Details** Target: WNT6 Alternative Name WNT6 (WNT6 Products) Background: Name/Gene ID: WNT6 Family: WNT Synonyms: WNT6, Protein Wnt-6 Gene ID: 7475 Pathways: WNT Signaling, Tube Formation **Application Details** Application Notes: Approved: IHC, IHC-P (15 µg/mL) Comment: Target Species of Antibody: Human Assay Procedure: The IHC-pro Immunohistochemistry Protocol **Tissue Preparation** Formalin fixation and embedding in paraffin wax **Tissue Sectioning** Make 4-µm sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-drying oven for 45 minutes at 60°C Deparaffinization Wash slides in 3 changes of xylene - 5 minutes each at room temperature. Rehydration Wash slides in 3 changes of 100% alcohol - 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol - 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block - 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water - 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene - 1 minute each at room temperature.

Apply coverslip

Restrictions:

For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-WNT6 antibody ABIN1049480 IHC staining of human breast, epithelium cells. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.